


Cell Host & Microbe

Deep mutational scanning of rabies glycoprotein defines mutational constraint and antibody-escape mutations

Graphical abstract



Authors

Arjun K. Aditham, Caelan E. Radford, Caleb R. Carr, Naveen Jasti, Neil P. King, Jesse D. Bloom

Correspondence

jbloom@fredhutch.org

In brief

Rabies virus causes thousands of human deaths annually and is the target of post-exposure prophylactic development. Aditham et al. use a pseudovirus platform to measure how all mutations to the rabies virus glycoprotein affect cell entry and antibody neutralization. This work will inform antibody therapy development and vaccine design.

Highlights

- A safe pseudovirus platform allows deep mutational scanning of rabies glycoprotein (G)
- Provide measurements detailing how all G mutations affect cell entry and antibody escape
- Identify sites likely involved in protein stability and conformational changes
- Identify naturally occurring mutations in rabies G that escape antibodies



Article

Deep mutational scanning of rabies glycoprotein defines mutational constraint and antibody-escape mutations

Arjun K. Aditham,¹ Caelan E. Radford,¹ Caleb R. Carr,^{2,3} Naveen Jasti,^{3,4,5} Neil P. King,^{4,5} and Jesse D. Bloom^{1,6,7,8,*}

¹Basic Sciences Division, Fred Hutchinson Cancer Center, Seattle, WA 98109, USA

²Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA

³Medical Scientist Training Program, University of Washington, Seattle, WA 98195, USA

⁴Department of Biochemistry, University of Washington, Seattle, WA 98195, USA

⁵Institute for Protein Design, University of Washington, Seattle, WA 98195, USA

⁶Computational Biology Division, Fred Hutchinson Cancer Center, Seattle, WA 98109, USA

⁷Howard Hughes Medical Institute, Seattle, WA 98109, USA

⁸Lead contact

*Correspondence: jbloom@fredhutch.org

<https://doi.org/10.1016/j.chom.2025.04.018>

SUMMARY

Rabies virus causes nearly 60,000 human deaths annually. Antibodies that target the rabies glycoprotein (G) are being developed as post-exposure prophylactics, but mutations in G can render such antibodies ineffective. Here, we use pseudovirus deep mutational scanning to measure how all single-amino-acid mutations to G affect cell entry and neutralization by a panel of antibodies. These measurements identify sites critical for G function and define constrained regions that are attractive epitopes for clinical antibodies, including at the apex and base of the protein. We provide complete maps of escape mutations for eight monoclonal antibodies, including some in clinical use or development. Escape mutations for most antibodies are present in some natural rabies strains. Overall, this work provides comprehensive information on the functional and antigenic effects of G mutations that can inform development of stabilized vaccine antigens and antibodies that are resilient to rabies genetic variation.

INTRODUCTION

Rabies is a lyssavirus that causes encephalitis that is fatal upon symptom onset in humans and most other mammalian species.¹ The virus is common in several mammalian species,^{2–4} and zoonotic human infections are responsible for ~60,000 human deaths annually, with the greatest toll in Africa and Asia.⁵ Exposure to rabies requires immediate post-exposure prophylaxis, which includes injections of rabies immunoglobulin (RIG) derived from human or equine serum, monoclonal antibodies, or repeated vaccination.⁶ Although prophylaxis is over 99% effective, cost and supply shortages render treatments inaccessible in some rabies-endemic regions.^{7–10} Therefore, efforts are underway to develop better rabies vaccines^{11–15} and monoclonal antibody treatments.^{16–19}

The main target of both vaccines and antibodies against rabies is its glycoprotein (G), which mediates receptor binding^{20–22} and viral fusion with host cells.^{23,24} Pre-fusion G is a trimeric complex on the viral surface,^{12,25} and undergoes pH-driven conformational changes to mediate membrane fusion.^{26,27} As for other class III viral fusion proteins,^{28–30} rabies G is conformationally dynamic, exchanging reversibly between its pre-fusion and post-

fusion conformations.^{24,31} G's conformational dynamics pose a challenge for vaccine design: the most potent neutralizing antibodies target the pre-fusion conformation, but many antibodies elicited by immunization with unmodified G protein bind to other conformations.^{12,25} Therefore, efforts are underway to stabilize pre-fusion rabies G to create improved vaccine antigens^{12,25} as well as identify potentially neutralizing anti-G monoclonal antibodies.^{19,32}

Another challenge is that G is quite diverse, with >10% protein sequence divergence among natural rabies strains. This genetic diversity complicates monoclonal antibody development, as natural strains have been identified with resistance to antibodies, including antibodies in clinical use or development.^{19,26,33–35} Attempts are being made to develop monoclonal antibodies or antibody cocktails with increased neutralization breadth and resistance to escape.^{19,34}

Here, we use pseudovirus deep mutational scanning^{36,37} to measure how mutations to the G ectodomain affect cell entry and neutralization by a panel of eight monoclonal antibodies. Our experiments quantify mutational constraint across G and provide insight into how mutations affect G's conformational dynamics and function. We also provide complete maps of escape






Figure 1. Deep mutational scanning of rabies G

(A) We create libraries of pseudoviruses expressing different mutants of G on their surface and encoding identifying nucleotide barcodes in their genomes. These libraries are used to infect cells in various conditions, and barcodes from viruses that successfully infect are deep sequenced to quantify the effects of mutations. (B) Schematic showing region of rabies G mutagenized in our libraries. We use the conventional numbering scheme where site 1 is assigned to the first site of ectodomain. In this numbering scheme, we mutagenized from site -2 to site 431.

(C) Effects of mutations on G-mediated cell entry relative to unmutated rabies G. Each column of the heatmap shows the effects of different amino acids at that site on cell entry, with the amino acid identity in the unmutated Pasteur strain G indicated with an "X." Mutations that impair cell entry G are colored orange, mutations that do not affect cell entry are white, and mutations that enhance cell entry are blue. Gray indicates mutations that were not reliably measured in the (legend continued on next page)

mutations for a variety of important antibodies, enabling us to better define their epitopes and assess the extent to which escape mutations for each antibody are present among natural rabies sequences. Overall, our deep mutational scan elucidates how mutations affect the function and antigenicity of G and provides information that can help guide the design of vaccine antigens and antibody treatments.

RESULTS

A pseudovirus deep mutational scan of rabies G

To measure the effects of mutations in rabies G, we used a previously described pseudovirus deep mutational scanning platform.^{36,37} This platform enables the creation of a library of genotype-to-phenotype linked lentiviral particles, each of which displays a unique G mutant on its surface and encodes an identifying nucleotide barcode in its genome (Figures S1A and S1B). These lentiviral particles (or “pseudoviruses”) encode only one viral gene (rabies G) and so can undergo only a single cycle of cellular infection, thereby enabling the study of G mutants without requiring the generation of replicative pathogenic virus. The pooled pseudovirus library can be used to infect cells under various experimental conditions, and the ability of each G mutant to enable cellular infection in each condition can be quantified by sequencing the barcodes from infected cells (Figure 1A). This approach makes it possible to measure the effects of thousands of different mutations to G in a single experiment.

We performed a deep mutational scan using the G protein from the Pasteur strain of rabies virus, a vaccine strain¹⁵ that recently had its pre-fusion G determined by cryoelectron microscopy (cryo-EM).^{12,25} We designed our mutant libraries to contain all single-amino-acid mutations to 433 sites in G, starting two residues N-terminal to the ectodomain (at the end of the signal peptide) and continuing through the entire structurally resolved^{12,25} region of the ectodomain and into its C-terminal region (Figure 1B). The libraries therefore target $433 \times 19 = 8,227$ different amino acid mutations. We also designed stop-codon mutations at 20 sites as negative controls for cell entry. Throughout this paper, we utilize the numbering scheme where one is assigned to the first site in the ectodomain, consistent with past numbering conventions for rabies G studies.^{12,25,26}

We created duplicate G pseudovirus mutant libraries using previously described methods (Figure S1B).^{36,37} Each library contained ~80,000 barcoded variants of G, and each covered >99% of the 8,227 intended ectodomain amino acid mutations (Figure S1C). Most barcoded variants contained only one G amino acid mutation (~64%), although some variants contained no (~12%) or multiple mutations (~24%) (Figure S1D). All sites were mutated at approximately equal frequency in each replicate library (Figure S1E), with the exception of site H384, which was missing most mutations (see STAR Methods). At most sites

except H384, all 19 possible amino acid mutations were observed in each replicate library (Figure S1F).

Effects of mutations on G-mediated cell entry

We first measured how all G mutations affected the protein’s ability to mediate pseudovirus entry into 293T cells. We did this by infecting 293T cells with pooled libraries of barcoded pseudovirus particles and miniprepping non-integrated reverse-transcribed lentiviral genomes.^{38–40} These lentiviral genomes are reverse transcribed into DNA only upon successful fusion of viral and cell membranes by the viral glycoprotein.^{41,42} We compared barcode frequencies in 293T cells infected with pseudovirus displaying the rabies G mutants against frequencies of each barcode in a control set of cells infected with barcoded pseudovirus that display unmutated vesicular stomatitis virus G protein (VSV-G) (Figures 1A, S1B, and S2A). To maximize titers of pseudovirus displaying rabies G, we performed infections with the G pseudoviruses in media supplemented with 4 μg/mL polybrene⁴³ and adjusted to a pH of 7.1⁴⁴ (Figure S2B). When pseudotyped with VSV-G, all pseudoviruses are able to enter 293T cells regardless of rabies G function, providing a baseline frequency for each barcode in the library. These baseline frequencies are then compared with the frequency of each barcode in the condition where the pseudoviruses are dependent on their rabies G mutant to infect cells.

We quantify the ability of each G variant to infect cells as the logarithm of the ratio of its barcode counts relative to those for the unmutated G between the rabies-G- and the VSV-G-mediated infection conditions (Figure S2A). Cell entry scores correspond to \log_2 changes in pseudovirus entry relative to the unmutated rabies G protein. Cell entry scores greater than zero indicate improved infection relative to parental G protein, whereas values below zero indicate impaired entry. Because a modest fraction of barcoded variants contain multiple mutations, we employed a global epistasis model to deconvolve the effects of individual mutations.⁴⁵ Measurements of how mutations affected cell entry were highly correlated between replicates performed with the same library on different days and between the two independent G mutant libraries (Figures S2C and S2D).

The effects of G mutations on cell entry are shown in Figure 1C; see the interactive displays at https://dms-vep.org/RABV_Pasteur_G_DMS/cell_entry.html to best examine the data. The effects of amino acid mutations on cell entry were bimodal: most mutations have either a neutral to slightly deleterious effect or a highly deleterious effect on cell entry (Figure S2E). As expected, mutations to stop codons are uniformly deleterious for cell entry. The effects of amino acid mutations vary widely across sites and, in some cases, among different mutations at the same site. For instance, sites 160–168 in the fusion domain, 181–189 in the third linker, and 201–206 and 244–250 in the pleckstrin homology domain were all tolerant of most amino acid mutations. But other sites are highly

deep mutational scanning. The overlay bar indicates the regions of G. See https://dms-vep.org/RABV_Pasteur_G_DMS/cell_entry.html for an interactive version of this heatmap.

(D) Validation of deep mutational scanning measurements of mutational effects on cell entry. For each of the 16 mutations (which were chosen to span a range of cell entry effects), the x axis shows the effect measured in the deep mutational scanning (DMS), whereas the y axis shows the titers of individual pseudoviruses generated with that G protein mutant relative to the unmutated parent. Two independent validation measurements of the titers were made for each mutation, hence there are two points per mutation in the plot. The *r* indicates the Pearson correlation.




Figure 2. Functional constraints in the context of rabies G's structure

(A) Pre-fusion G structure (PDB: 7U9G) with two transparent protomers and one protomer colored by mean effect of all mutations at each site on cell entry. Sites are colored orange when mutations impair cell entry and white when they have no effect. See https://dms-vep.org/RABV_Pasteur_G_DMS/cell_entry.html#mutation-effects-on-structure-of-g for interactive structural visualizations of the effects of mutations.

(B) Zoomed view of the fusion domain of a single protomer of pre-fusion G with sites colored by the mean effects of mutations on cell entry. The fusion loops are not resolved in the structure and are modeled as dashed cartoon loops.

(C) Heatmaps showing the effects on cell entry of all mutations in the fusion loops and flanking sites. The parental amino acid identities in the Pasteur strain G are indicated with "X."

(D) Zoomed view of a single protomer of pre-fusion G highlighting the histidine cluster and structurally adjacent threonines that may act as a pH sensor. Sites are colored by the mean effect mutations on cell entry as in (A). See Figure S4 for validation of effects of some mutations in the histidine cluster.

(E) A single protomer of G shown in both the pre-fusion protomer and extended intermediate conformations (PDB: 6LGW), with sites highlighted in (F) shown in red and sites highlighted in (G) shown in blue.

(legend continued on next page)

constrained: for instance, nearly all mutations are deleterious in both fusion loops, which span sites 73–79 and 117–125.²⁶ In the next subsection, we discuss these patterns of constraint in more detail in the context of G's structure.

To validate the deep mutational scanning measurements of mutation effects on cell entry, we cloned 15 individual G mutations measured to have a range of effects and determined the titers of individual pseudoviruses encoding these mutations. The effects of mutations on cell entry from the deep mutational scan were highly correlated with the individual pseudovirus titers measured in these validation assays ($r = 0.91$; Figure 1D). The deep mutational scanning is also consistent with prior experiments that have characterized the effects of individual mutations to G on its fusogenic activity^{12,26} (Figure S2F).

Mutational constraint in the context of G's structure and function

The effects of mutations on G's cell-entry function varied widely across the protein, with both highly constrained and mutationally tolerant sites in all three of the central domains, pleckstrin homology domain, and fusion domain (Figures 1C and 2A). Some of this constraint can be explained relatively simply by the requirement that G fold properly in order to function. For example, one of the most constrained regions on G's surface involves the disulfide bond connecting sites C189 and C228, where mutations to any non-cysteine amino acid at either position are highly deleterious for cell entry (Figure 2A). At sites that are buried in G's pre-fusion structure, mutations to non-hydrophobic amino acids tend to be deleterious, presumably because they disrupt the folding of the protein's hydrophobic cores (Figure S3).

Other sites of constraint provide more nuanced insights into G's function. The fusion loops are critical to G's cell entry mechanism and are thought to stabilize the trimeric pre-fusion conformation of rabies G.²⁵ The fusion loops are only partially resolved in available three-dimensional structures of G, but our results show that many sites in both the loops and flanking beta strands are highly constrained (Figures 2B and 2C). Prior work has tested alanine mutations to sites F74, Y77, Y119, and W121 in the fusion loops on stability of trimeric G and found that only Y119A does not have a strong adverse effect.²⁵ Consistent with this prior work, our deep mutational scan shows that a mutation to alanine is reasonably tolerated for cell entry at Y119 but not at F74, Y77, or W121 (Figure 2C). However, all of these sites except W121 do tolerate mutations to some amino acids that conserve hydrophobicity or aromaticity (e.g., F74L and Y77F are reasonably tolerated) (Figure 2C). Proline residues near the fusion loops (sites 89, 106, 115, and 117) are largely intolerant of mutations (Figures 1C and 2B), likely because these prolines are important for ensuring proper fusion loop orientation.

The deep mutational scanning also helps delineate sites that may be important in triggering the conformational change in G necessary for cell entry. A cluster of histidines at sites 86, 173, and 397 at the interface between one of the linkers and the fusion domain is hypothesized to help govern pH-driven conforma-

tional change in rabies G^{12,46} (Figure 2D). Consistent with this hypothesis that these histidines act as a "pH sensor" crucial for G's cell-entry function, our deep mutational scan shows that mutating any of these three sites to any non-histidine amino acid impairs cell entry (Figures 1C and 2D). We validated that the mutations H86A and H397A impair G's cell-entry function without reducing cell-surface expression of pre-fusion G, consistent with the idea that these mutations impair the pH sensor that activates G for fusion (Figure S4; note that H173A impairs G surface level expression). Our deep mutational scan also shows that threonine residues structurally adjacent to the histidines (T60, T62, T171, and T176) only tolerate serine mutations (Figures 1C and 2D), suggesting that the hydroxyl moieties of these residues may be important for G function. This finding is consistent with the possibility that these residues interact with the histidine cluster, as has been hypothesized for serines near the histidine cluster in the related VSV-G fusion protein.⁴⁶

Our deep mutational scanning identifies some sites where the mutational constraint appears to be due to interactions in the extended intermediate conformation, which exists in equilibrium with the pre-fusion trimer^{24,27,31} (Figure 2E). A cluster of sites that includes E45, D211, R213, H261, D266, and E269 form polar and van der Waals interactions in the extended intermediate¹² (Figure 2F), and our deep mutational scanning shows that almost all mutations at these sites are deleterious for cell entry (Figures 1C and 2F). Another cluster of sites (L46, P144, Y145, I191, F192, L227, W240, and V258) form extensive hydrophobic contacts in the extended intermediate (Figures 2E and 2F), and our deep mutational scanning shows that only hydrophobic mutations (or in the case of Y145, no mutations) are tolerated at these sites (Figures 1C and 2F). In the pre-fusion conformation trimer, some of these sites (e.g., Y145 and V258) are distal from the other sites and largely solvent exposed (Figure 2E), suggesting that the mutational constraints originate from interactions in the extended intermediate, rather than the pre-fusion conformation. These sites that stabilize the extended intermediate but likely tolerate mutation in the pre-fusion conformation represent additional candidates for stabilizing mutations that might shift the conformational equilibrium of rabies G toward the pre-fusion state.

Prior work has stabilized rabies G in its pre-fusion conformation by introducing a mutation to proline (H270P) that is tolerated in the pre-fusion conformation but is disfavored in the extended intermediate conformation.^{12,13} A similar proline-based stabilization strategy has been applied to stabilize vaccine immunogens for both class I fusion proteins (e.g., coronavirus spikes^{47,48} and Lassa glycoprotein complex⁴⁹) and VSV-G.⁵⁰ Consistent with the idea that a proline at site 270 in rabies G uniquely blocks conversion of the pre-fusion trimer to the extended intermediate, our deep mutational scanning shows that H270P strongly impairs cell entry but most other amino acid mutations at this site are well tolerated (Figure S5). We used our deep mutational scanning to identify additional sites in regions that undergo large conformation changes where proline is especially disfavored

(F) Zoomed view of the extended intermediate conformation showing sites that form polar interactions in this conformation. Sites are colored by the mean effects of mutations on cell entry as in (A).

(G) Zoomed view of the extended intermediate conformation showing sites that form hydrophobic interactions in this conformation. Sites are colored by the mean effects of mutations on cell entry as in (A).




Figure 3. Deep mutational scanning of how G mutations affect antibody neutralization, as applied to antibody 17C7

(A) Workflow for mapping effects of mutations on antibody neutralization. The rabies G pseudovirus library is mixed with a “non-neutralized standard” consisting of barcoded VSV-G pseudovirus, then this pool is incubated with different concentrations of antibody and used to infect 293T cells. Barcodes are recovered from infected cells and deep sequenced to quantify the ability of each variant to infect cells at each antibody concentration. The VSV-G non-neutralized standard pseudovirus is not affected by anti-rabies G antibodies and so is used to convert the sequencing counts into the fraction of each variant that is neutralized at each antibody concentration.

(B) Line plot showing the total escape caused by mutations at each site in G for antibody 17C7. See https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html for interactive visualizations of escape. Orange marks near the x axis indicate sites zoomed in the logo plot in (C).

(C) Logo plot of key sites of escape from antibody 17C7. The height of each letter is proportional to the escape caused by that amino acid mutation. Total height of letters at each site is proportional to the sum of all mutant escape scores. The letters are colored according to the effect of each mutation on cell entry.

(D) Structure of 17C7 and rabies G (PDB: 8A1E), with G colored by the total escape caused by mutations at each site (blue indicates sites where mutations cause the most escape). The antibody is shown in purple.

(E) Validation assays comparing the escape measured in the DMS versus the IC50s measured in standard pseudovirus neutralization assays against antibody 17C7 for nine different single-amino-acid mutants of G. Raw neutralization curves are in Figure S5. The mutations tested in these validation assays were chosen to span a range of escape values as measured in the DMS. Pseudovirus with unmutated rabies G was measured in triplicate, and mean \pm standard deviation is plotted.

(e.g., K47, Q256, N259, L260, and S265) (Figures S5A–S5C). We validated that mutations Q256P, L260P, and S265P (as well as the previously known H270P mutation) all impair cell entry without reducing cell-surface expression of pre-fusion G, consistent with the hypothesis that they lock the protein into the pre-fusion conformation (Figures S5D and S5E). Therefore, these sites represent additional potential candidates for pre-fusion stabilizing proline mutations for immunogen design.

Deep mutational scanning comprehensively maps antibody-escape mutations

To experimentally measure how all rabies G mutations affect antibody neutralization, we incubated the pseudovirus library

with different concentrations of antibody and used deep sequencing to quantify the ability of each G variant to infect 293T cells at each antibody concentration (Figure 3A). To convert the deep-sequencing counts to neutralization values, we normalized them to the counts of a “non-neutralized standard” consisting of VSV-G pseudovirus that is not neutralized by the anti-rabies antibodies³⁶ (Figure 3A). We analyzed the data using a previously described biophysical model⁵¹ to obtain escape values for each mutation that are proportional to the log fold change in IC50 (half-maximal inhibitory concentration).

We first applied this approach to monoclonal antibody 17C7, which is the active component of the Rabishield post-exposure prophylactic^{16,33,52} and has had its structure in complex with

Table 1. Antibodies with escape mutations mapped in this study

Antibody	Development status	Antigenic region	PDB ID	Pan-lyssavirus activity ³⁴	IC50 (ng/mL)
17C7 ¹⁶	clinically deployed in India as of 2017 ^{19,57}	III ³⁴	8A1E ¹²	no	0.35
CR4098 ⁵⁵	part of CL184 cocktail ^{55,61} ; development stopped after phase II trial ^{19,32}	III ⁶²	N/A	no	3.8
RVA122 ³⁴	N/A	III ³⁴	7U9G ²⁵	yes	0.31
RVC58 ³⁴	animal testing in cocktail with RVC20 ⁶³	III ³⁴	N/A	yes	0.22
RVC20 ³⁴	animal testing in cocktail with RVC58 ⁶³	I ³⁴	6TOU ⁵⁶	yes	0.36
CR57 ⁶⁴	part of CL184 cocktail (development stopped after phase II trial) ^{19,32} ; part of cocktails in strain panel testing ^{19,33}	I ⁶⁴	8R40 ⁶⁵	yes	0.55
CTB012 ¹⁸	part of SYN023 cocktail; clinical trials ongoing ^{58,59} (phase III complete ⁶⁰)	N/A ^{18,35}	N/A	untested	0.47
RVC68 ³⁴	N/A	N/A ³⁴	N/A	yes	7.6

IC50s are from pseudovirus neutralization assays (Figure S5B). Antigenic regions were taken from De Benedictis et al.³⁴ and de Melo et al.^{19,63} and antibodies marked “N/A” bind outside known antigenic regions. Pan-lyssavirus activity was defined based on strain panel testing from De Benedictis et al.³⁴

G determined by cryo-EM.¹² Our deep mutational scanning showed that mutations that escape neutralization by 17C7 occur predominantly at sites 334, 336–338, 340–342, and 346 (Figures 3B and 3C and interactive visualizations at https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html). We visualized the escape caused by mutations at key sites in logo plots where the height of each amino acid letter is proportional to that mutant’s escape score, and the overall height at each site is proportional to the sum of all mutant escape scores (Figure 3C). Many of these escape mutations are well tolerated with respect to G’s cell-entry function (as shown by the color of letters in Figure 3C). The escape mutations all occur near the binding interface of 17C7 and G (Figure 3D). One of the escape mutations, I338T, does not directly contact the antibody but creates an N-linked glycosylation motif at N336 that is known to affect G’s antigenicity.^{53,54} I338S does not appear as an escape mutation because, unlike I338T, it is highly deleterious for cell entry in the background of the Pasteur strain G (Figure 1C).

To validate the deep mutational scanning measurements of antibody escape, we created individual pseudoviruses encoding nine different mutations that had a range of effects on neutralization by antibody 17C7 in the deep mutational scanning. The effects of these mutations, as measured in traditional neutralization assays performed with these pseudoviruses, were highly correlated with the escape values measured in the deep mutational scanning ($r = 0.97$; Figures 3E and S6A), validating the accuracy of the approach.

Comprehensive escape maps for eight monoclonal antibodies

We used deep mutational scanning to experimentally map all escape mutations for eight monoclonal antibodies targeting G (Table 1). These antibodies bind to different antigenic regions on G: two target antigenic region I (antibodies CR57^{54,55} and RVC20^{34,56}), four target region III (17C7,¹⁶ CR4098,⁵⁵ RVA122,^{25,34} and RVC58^{17,34}), and two bind outside previously described antigenic regions (CTB012¹⁸ and RVC68³⁴). Antibody 17C7 has been clinically deployed in India since 2017,^{19,57} and

CTB012¹⁸ is one of the two active components of the SYN023 cocktail^{58,59} that recently concluded phase III clinical trials.⁶⁰ All eight antibodies neutralized pseudovirus with unmutated rabies G, albeit with varying potencies (Table 1; Figure S6B).

The deep mutational scanning shows that each antibody is escaped by mutations at a relatively small number of sites that spatially cluster in the pre-fusion structure of G (Figure 4 and interactive visualizations at https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html). For the four antibodies that have had their structures in complex with G experimentally determined, the major escape mutations all occur at sites in G near the antibody-binding interface (Figures 5A and 5B). However, mutations at only a fraction of the contact sites strongly escape antibody neutralization (Figures 5A and 5B), a result consistent with prior work on antibody escape for other viruses.^{66–69} In some cases, certain contact sites have no escape mutations simply because all mutations are too deleterious for G’s cell-entry function (e.g., site C228 for antibody RVC20 in Figure 5B), but, in other cases, contact sites tolerate many mutations but none cause appreciable escape from neutralization (e.g., site E33 for antibody RVA122 in Figures 5A and 5B).

Antibodies that target the same antigenic region of G can differ substantially in their escape mutations. For instance, 17C7, CR4098, RVA122, and RVC58 all bind to antigenic region III of G.³⁴ However, although 17C7 and CR4098 are escaped by mutations at very similar sets of sites, the sites of escape mutations for RVA122 and RVC58 are more distinct (Figure 4). Even when antibodies share sites of escape, the effects of individual mutations can differ. For instance, N336S³⁵ and R346K^{70,71} escape 17C7 but not CR4098, although some other mutations at sites 336 and 346 escape both antibodies (Figure 4). Likewise, RVC20 and CR57 both bind to antigenic region I,³⁴ where they are believed to block the binding site for the alpha7 nicotinic acetylcholine receptor^{22,65,72} and prevent pH-driven conformational change.^{56,65} Despite the similarity of their epitopes, the escape mutations for these two antibodies only partially overlap (Figures 4, 5A, and 5B). For instance, both antibodies are affected by mutations to K226 and G229, but mutations at V230 and A242 only strongly escape CR57.




Figure 4. Comprehensive maps of escape mutations for eight monoclonal antibodies

For each antibody, the line plot shows the total escape caused by all mutations at that site; orange marks near the x axis indicate sites zoomed in the logo plots. The logo plots show the escape caused by individual mutations at key sites, with letter heights proportional to escape value for mutation to that amino acid and

(legend continued on next page)

The escape maps also provide insight into why some antibodies have greater breadth against lyssaviruses than others. For instance, among the antibodies targeting antigenic region III, 17C7 and CR4098 only neutralize rabies, whereas RVA122 and RVC58 also broadly neutralize non-rabies lyssaviruses (e.g., Duvenhage lyssavirus, European bat 1 lyssavirus, and Australian bat lyssavirus).³⁴ Unlike the narrow-spectrum antibodies (17C7 and CR4098), the broad-spectrum antibodies (RVA122 and RVC58) are escaped by mutations at fewer sites or sites with lower mutational tolerance. Specifically, RVA122 is mostly escaped by mutations at just one site: R333 (Figures 4, 5A, and 5B), a site that is commonly mutated in vaccine strains^{73–75} but where mutations generally attenuate pathogenicity.^{76,77} RVC58 is escaped by mutations at several sites, but many of those mutations, such as those at sites L38 and V210, are deleterious to G's cell-entry function (Figure 4, see color of letters in logo plots).

Two of the antibodies (CTB102 and RVC68) lack structures in complex with G and do not bind to previously defined antigenic regions, although the epitope of CTB102 has been partially defined by alanine mutagenesis and escape studies.¹⁸ Our deep mutational scanning shows that CTB102 binds to sites on the apex of G that are spatially clustered on pre-fusion rabies G, primarily in the loops involving V29-E33 and I268-L275 (Figures 4 and 5C). These loops undergo a substantial rearrangement in the extended intermediate conformation of G,^{24,27,31} meaning that the epitope of CTB102 is only spatially contiguous in the pre-fusion conformation of G (Figure 5C). RVC68 has very broad neutralization activity across lyssaviruses,³⁴ but relatively poor neutralization potency against rabies G (Table 1). Our deep mutational scanning shows that RVC68 targets the base of the fusion domain of rabies G (Figure 4). The broad activity of this antibody against different lyssaviruses likely stems from conservation of the targeted sites in previously tested viruses.³⁴

Antibody-escape mutations are present in naturally occurring rabies strains

The G protein varies among natural rabies strains, and some strains have been identified that are resistant to specific antibodies, including clinically relevant antibodies.^{33,34,54,70} Our deep mutational scanning enables systematic identification of antibody-resistant strains. We first examined the frequency of each amino acid mutation among naturally occurring G sequences versus its antibody escape measured in our deep mutational scan (Figure 6A and interactive plot at https://dms-vep.org/RABV_Pasteur_G_DMS/natural_seqs.html). Mutations that cause at least some escape are present among natural G sequences for all antibodies, but the frequencies of these mutations and the magnitude of escape they cause differ markedly among antibodies. For example, many strong escape mutations from antibody 17C7 are at appreciable frequencies among natural sequences (Figure 6A). In contrast, antibodies such as RVC20 have only a few modest-effect escape mutations at appreciable frequency among natural sequences (Figure 6A).

We predicted each rabies strain's escape from each antibody by summing the experimentally measured effects of all of its constituent G mutations relative to the Pasteur strain. We visualized these predictions via Nextstrain⁷⁸ phylogenetic trees, where sequences can be colored by their predicted escape (<https://nextstrain.org/groups/jbloombloom/dms/rabies-G>). As an example, a static view of the tree for all 7,122 publicly available G sequences colored by escape from antibody 17C7 is shown in Figure 6B. Although most strains lack escape mutations from 17C7 (white in Figure 6B), there are some strains predicted to strongly escape for this antibody (blue in Figure 6B). Most of these strains contain mutations at sites 336 or 346 (Figure S7A), both of which are sites where mutations are measured to strongly escape 17C7 in the deep mutational scan (Figures 4, 6A, and 6C). Note that many of the sequenced strains that are predicted to most strongly escape 17C7 originate from infected bats or wild animals rather than dogs (Figure 6B), which are the most common source of human infections.⁷⁹

As validation, we identified two natural strains predicted to strongly escape antibody 17C7, NY-2011-12020 and A12_2718 (Figure 6C). The G proteins of these two strains are distantly related to the Pasteur strain, as both differ from it by at least 35 amino acid mutations. We generated pseudovirus expressing G from each of these strains and measured neutralization by 17C7 as well as another antibody predicted not to be escaped by each of the strains (Figures 6C and 6D). As predicted by the deep mutational scanning, both strains strongly escaped neutralization by 17C7 but retained neutralization by the control antibodies at levels similar to the Pasteur strain (Figure 6D).

To test how much the escape was explained by just the top escape mutations identified in the deep mutational scanning, we either removed each of the two top escape mutations (I338T or R346S) from the A12_2718 strain G or introduced them into the Pasteur strain G. Consistent with the deep mutational scan, removing the top escape mutation of I338T from the A12_2718 G partially restored neutralization by 17C7, whereas adding this single mutation to the Pasteur G made it fully resistant to neutralization by 17C7 (Figure S7B). Introducing the weaker escape mutation R346S to the Pasteur strain reduced but did not eliminate its neutralization by 17C7, also in agreement with the deep mutational scan (Figure S7B). Although these measurements validate the ability of deep mutational scanning to predict escape of substantially divergent strains for the antibodies tested here, caution should always be exercised when extrapolating the deep mutational scanning to sequence-divergent strains. In some cases, epistasis may lead to shifts in the effects of mutations, as observed in other viral proteins.^{80–82}

DISCUSSION

Here, we used deep mutational scanning to measure how mutations to rabies G affect cell entry and antibody neutralization. The comprehensive measurements of how mutations affect

letters colored according to the effects of mutations on cell entry (see color scale at the bottom right of the figure). The structures show pre-fusion rabies G (PDB: 7U9G), with residues colored by the total escape caused by mutations at each site (see color scale at bottom right of the figure). See Figure 5 for additional structural renderings of antibodies with experimentally solved structures in complex with G. See https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html for interactive heatmaps and structure-based visualizations of escape.




Figure 5. Mutations at only some of the antibody contact sites in G mediate escape

(A) G colored by extent of antibody escape at each site, as measured in deep mutational scanning for the four antibodies with experimentally determined structures in complex with G (PDB: 8A1E, PDB: 7U9G, PDB: 6TOU, and PDB: 8R40). The antibodies are shown in magenta. The color scale for G is defined at the lower left of the panel: white indicates no escape at the site, and blue indicates strong escape at the site.

(B) Escape values for each mutation for all sites in G that contact each antibody in the experimentally determined structural complex. A site in G is defined as being in contact if any atom is within 4 Å of the antibody. Mutations are colored white to blue, based on the escape values measured in deep mutational scanning. Mutations that are too deleterious for cell entry to reliably measure their effect on antibody neutralization are shown in dark gray. The small number of mutations in light gray (e.g., Q382F and K226H) have no measured escape due to insufficient coverage in the library. The amino acid identity in the unmutated Pasteur G is indicated with an "X" for each site.

(C) Structures of rabies G in pre-fusion (PDB: 7U9G) and extended intermediate (PDB: 6LGW) conformations colored with sites colored by the total escape from the CTB012 antibody. Escape sites are contiguous in pre-fusion rabies G but more distant in extended intermediate conformation. For visibility, key sites are shown in spheres in the extended intermediate. Color scale is the same as for (A).




Figure 6. Naturally occurring antibody-escape mutations in rabies G

(A) Frequency of each G amino acid mutation among naturally occurring rabies sequences versus the escape caused by that mutation in the deep mutational scanning of the Pasteur strain G. Each panel shows escape from a different antibody. Dashed vertical lines indicate mutations observed at least two times in natural strains; points in light gray indicate rare mutations. Mutations not observed in natural strains are assigned a frequency of 10^{-5} to enable plotting on a log scale. See https://dms-vep.org/RABV_Pasteur_G_DMS/natural_seqs.html for an interactive version of this plot that allows mouseover of individual points to see the mutation identities.

(B) Phylogenetic tree of all publicly available 7,122 rabies G sequences colored by escape from antibody 17C7, as predicted by summing the experimentally measured effects of all mutations relative to the Pasteur strain. White indicates no escape and blue indicates strong escape. See <https://nextstrain.org/groups/bloomlab/dms/rabies-G> for interactive Nextstrain phylogenies showing the predicted escape of all strains for each antibody. The tree is rooted by Gannoruwa bat lyssavirus (accession: NC_031988.1), a non-rabies lyssavirus also in phylogroup I. Strains experimentally tested in validation neutralization assays are labeled. Common hosts for different clades are labeled. The interactive Nextstrain phylogenies provide the option to label all strains by host.

(C) For validation neutralization assays, we identified antibodies predicted to be escaped or neutralized by two different natural strains. The table summarizes the strains and antibodies used in the validation assays. The numbers in parentheses after the antibody names and mutations in the last two columns are the escape values from the deep mutational scanning.

(D) Neutralization curves of pseudovirus expressing G from the Pasteur, NY-2011-12020, or A12_2718 strains against the indicated antibodies. Points indicate the mean \pm standard error of technical duplicates.

G's cell-entry function define which sites in the protein have the most capacity for evolutionary change and shed light on how mutations affect G's conformational dynamics in a way that could inform design of vaccine antigens. The measurements of how mutations affect antibody neutralization delineate key epitopes. This information can guide the development of antibodies and antibody cocktails that are robust to G's natural genetic diversity, as well as define key epitopes to preserve in vaccine design.

With respect to how mutations affect G's cell-entry function, we find that some regions of G (e.g., the base of the fusion domain and parts of the trimer apex) are intolerant of most mutations, but other regions can tolerate mutations without substantially impairing cell entry. In some instances, we can identify sites where mutational constraint appears to arise from protein contacts formed in just one of the pre-fusion or extended intermediate conformations. Mutations that differentially affect these two conformations could aid in the design of vaccine immuno-

gens. The conformational dynamics of rabies G are hypothesized to hamper the neutralizing antibody response to vaccination by unmodified G because some potentially neutralizing antibodies target only pre-fusion G.^{13,25,26} Several studies have therefore attempted to stabilize pre-fusion G by introducing proline mutations that block conformational changes,^{12,13} similar to what has been done for other viral fusion proteins.^{47,48,50} Our work suggests an additional mechanism of pre-fusion stabilization by identifying sites where mutations may destabilize the extended intermediate.

Our work better defines how neutralizing antibodies target G and how these antibodies are affected by viral mutations. Delineation of which viral mutations escape anti-G antibodies is important because monoclonal antibodies are being developed for post-exposure prophylaxis, with some already in clinical use. We comprehensively mapped escape mutations for eight antibodies, including the antibody in Rabishield and several other antibodies under development for use against rabies.^{16–18,63} Some

of these antibodies have had their structures in complex with G previously determined, and our work shows that only some mutations at the antibody-G interface confer escape. Other antibodies that we mapped have not been structurally characterized, and, for those antibodies, our work defines their epitopes. Furthermore, our measurements help rationalize why antibodies differ in their breadth against lyssaviruses: the broadest antibodies tend to target sites in G where most mutations are detrimental to G's cell-entry function.

Our deep mutational scanning can help identify which natural rabies strains are likely to escape specific antibodies. Strikingly, we find that escape mutations to all eight antibodies we examined are present among natural rabies strains—but escape mutations are far more common for some antibodies than others. We provide interactive phylogenetic trees that predict the escape of publicly available rabies G sequences based on the effects of their constituent mutations as measured in our experiments. These predictions are approximate because there can be epistasis among combinations of mutations^{80–82} (see the [limitations of the study](#) section below). However, we did experimentally validate several examples showing that our data can predict the susceptibility or resistance of natural strains to specific antibodies. Going forward, the deep mutational scanning data provide a powerful way to identify natural strains that are likely to harbor resistance mutations, although such predictions should always be independently validated given the potential for epistasis.

Overall, our work provides the first large-scale measurements of the functional and antigenic effects of mutations to rabies G. Our data can be interrogated in the context of G's structure and evolution via interactive visualizations we have made publicly available (https://dms-vep.org/RABV_Pasteur_G_DMS/), which should facilitate their use in basic research, vaccine and antibody development, and surveillance of rabies virus evolution.

Limitations of the study

Our experiments measure how mutations to G affect its ability to mediate pseudovirus infection of 293T cells. However, pseudovirus infection of 293T cells does not recapitulate all aspects of actual rabies virus infection *in vivo*.^{3,83,84} Therefore, although our experiments do capture the core processes of G-mediated cell entry and antibody neutralization, they may not authentically reflect some more subtle aspects of real viral infection, such as differential usage of specific receptors.^{20,72,85} Cell entry also reflects the combined effects of biochemical perturbations to several properties of G, such as expression level, stability, and pH-dependent activation.

Our deep mutational scanning used G from the lab-adapted Pasteur vaccine strain, which has been passaged extensively in animals and cell lines.⁸⁶ Some of the mutations from this passaging enhance pseudovirus production^{87,88} but could alter some of G's properties relative to natural strains. Additionally, there is always the potential for epistasis among mutations,^{80–82} such that the effect of a mutation in the Pasteur G could differ for glycoproteins from other strains with multiple mutations. This limitation is especially relevant when using deep mutational scanning to predict the antibody escape of other rabies strains (last section of [results](#)). Although such predictions validated well for distantly related rabies G sequences

we tested here, caution should always be exercised when predicting properties of divergent strains from measurements made on the Pasteur G.

Finally, like all high-throughput experiments, deep mutational scanning is subject to experimental noise. We performed all experiments in replicate using independently generated pseudovirus libraries, but, in most of this paper, we simply report the average of these replicates. For mutations of special interest, we recommend also examining the interactive heatmaps at https://dms-vep.org/RABV_Pasteur_G_DMS/, where you can mouseover points to look at the measurements for each individual replicate and view other quality control metrics (such as the number of unique variants in which each mutation was observed).

RESOURCE AVAILABILITY

Lead contact

Requests for reagents and resources should be directed to the lead contact, Jesse D. Bloom (jbloom@fredhutch.org).

Materials availability

Materials associated with this study are available upon request. All unique reagents generated in this study are available from the [lead contact](#) without restriction.

Data and code availability

See https://dms-vep.org/RABV_Pasteur_G_DMS/ for links to interactive visualizations of the data and CSVs containing the final inferred mutational results, including structure-based visualizations rendered with *dms-viz*.⁸⁹ See https://github.com/dms-vep/RABV_Pasteur_G_DMS for the full computer and all data starting with the barcode counts.

The interactive plots at https://dms-vep.org/RABV_Pasteur_G_DMS/ contain a number of options that enable different displays of the data or filtering by the confidence of the measurement. In the heatmaps, you can mouseover points for details on the per-replicate measurements (note all mutations were measured in at least two libraries). There are also options to adjust the site summary statistic (e.g., mean or sum of mutation effects at a site), zoom in on particular regions, and filter values as follows:

- minimum times_seen: average number of variants per-library that contain a specific mutation; mutations present in more variants are generally measured more accurately.
- maximum effect_std: maximum standard deviation of measurement across all replicates.
- minimum n_selections or minimum n_models: minimum number of replicates in which a particular mutation was measured.
- minimum max of effect at site or minimum max escape at site: only show sites for which at least one mutation has a measured value greater than this.
- minimum functional effect: for antibody-escape, gray out and ignore any mutations with an effect worse than this on cell entry; this filters highly deleterious mutations.

These filters are all set to reasonable values by default, but, in some cases, you may want to adjust them further. In the heatmaps, mutations that lack measurements that meet the above filtering criteria are shown in light gray; in the antibody-escape heatmaps, mutations that are highly deleterious are shown in dark gray.

ACKNOWLEDGMENTS

We thank Brendan Larsen, William Hannon, and Sophie Shoemaker for technical advice and feedback. This work was supported by NIH grant R01 AI141707 to J.D.B. and ARPA-H contract 1AY1AX000036-01 to J.D.B. and N.P.K. We thank the Genomics & Bioinformatics Shared Resource

(RRID: SCR_022606) of the Fred Hutchinson Cancer Center (P30 CA015704). This research was supported by the Flow Cytometry Shared Resource, RRID: SCR_022613, of the Fred Hutch/University of Washington/Seattle Children's Cancer Consortium (P30 CA015704). J.D.B. is an investigator of the Howard Hughes Medical Institute.

AUTHOR CONTRIBUTIONS

Conceptualization, A.K.A. and J.D.B.; methodology, A.K.A., C.E.R., and J.D.B.; software, C.E.R., C.R.C., and J.D.B.; investigation, A.K.A.; resources, N.J. and N.P.K.; data curation, A.K.A., C.E.R., C.R.C., and J.D.B.; visualization, A.K.A., C.R.C., and J.D.B.; writing – original draft, A.K.A. and J.D.B.; writing – review and editing, A.K.A., C.E.R., C.R.C., N.J., N.P.K., and J.D.B.; supervision, J.D.B.; funding acquisition, J.D.B. and N.P.K.

DECLARATION OF INTERESTS

J.D.B. consults on topics related to viral evolution for Apriori Bio, Invivyd, the Vaccine Company, and Pfizer. J.D.B., A.K.A., and C.E.R. are inventors on Fred Hutch licensed patents related to viral deep mutational scanning. N.P.K. consults on topics related to vaccine design for AstraZeneca, and the King lab has received unrelated sponsored research agreements from Pfizer and GlaxoSmithKline.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **EXPERIMENTAL MODEL DETAILS**
 - Rabies G used for deep mutational scanning
 - Cells
 - Antibodies
- **METHOD DETAILS**
 - G plasmid mutant libraries
 - Cell-stored rabies G mutant library
 - Production of rabies G and VSV-G expressing pseudovirus libraries
 - Long-read sequencing to link barcodes with rabies G mutants
 - Deep mutational scanning for mutational effects on cell entry
 - Deep mutational scanning for mutational effects on antibody escape
 - Illumina sequencing of pseudovirus barcodes
 - Production of luciferase-encoding pseudovirus for validation assays
 - Validation pseudovirus neutralization assays
 - Pseudovirus validations of mutational effects on cell entry and antibody escape
 - Validation of deep mutational scanning predictions using natural rabies G strains
 - Measuring cell surface expression levels of rabies G mutants
 - Computational Methods
- **QUANTIFICATION AND STATISTICAL ANALYSIS**

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.chom.2025.04.018>.

Received: December 18, 2024
Revised: March 25, 2025
Accepted: April 23, 2025
Published: May 20, 2025

REFERENCES

1. Fooks, A.R., Banyard, A.C., Horton, D.L., Johnson, N., McElhinney, L.M., and Jackson, A.C. (2014). Current status of rabies and prospects for elimination. *Lancet* 384, 1389–1399. [https://doi.org/10.1016/S0140-6736\(13\)62707-5](https://doi.org/10.1016/S0140-6736(13)62707-5).
2. Mollentze, N., Biek, R., and Streicker, D.G. (2014). The role of viral evolution in rabies host shifts and emergence. *Curr. Opin. Virol.* 8, 68–72. <https://doi.org/10.1016/j.coviro.2014.07.004>.
3. Fisher, C.R., Streicker, D.G., and Schnell, M.J. (2018). The spread and evolution of rabies virus: conquering new frontiers. *Nat. Rev. Microbiol.* 16, 241–255. <https://doi.org/10.1038/nrmicro.2018.11>.
4. Singh, R., Singh, K.P., Cherian, S., Saminathan, M., Kapoor, S., Manjunatha Reddy, G.B., Panda, S., and Dhama, K. (2017). Rabies-epidemiology, pathogenesis, public health concerns and advances in diagnosis and control: a comprehensive review. *Vet. Q.* 37, 212–251. <https://doi.org/10.1080/01652176.2017.1343516>.
5. Hampson, K., Coudeville, L., Lembo, T., Sambo, M., Kieffer, A., Attlan, M., Barrat, J., Blanton, J.D., Briggs, D.J., Cleaveland, S., et al. (2015). Estimating the Global Burden of Endemic Canine Rabies. *PLoS Negl. Trop. Dis.* 9, e0003709. <https://doi.org/10.1371/journal.pntd.0003709>.
6. Nadal, D., Bote, K., Masthi, R., Narayana, A., Ross, Y., Wallace, R., and Abela, B. (2023). Rabies post-exposure prophylaxis delivery to ensure treatment efficacy and increase compliance. *IJID One Health* 1, 100006. <https://doi.org/10.1016/j.ijidoh.2023.100006>.
7. Chungalucha, J., Steenson, R., Grieve, E., Cleaveland, S., Lembo, T., Lushasi, K., Mchau, G., Mtema, Z., Sambo, M., Nanai, A., et al. (2019). The need to improve access to rabies post-exposure vaccines: Lessons from Tanzania. *Vaccine* 37, A45–A53. <https://doi.org/10.1016/j.vaccine.2018.08.086>.
8. World Health Organization (2018). Rabies vaccines: WHO position paper, April 2018 – Recommendations. *Vaccine* 36, 5500–5503. <https://doi.org/10.1016/j.vaccine.2018.06.061>.
9. Wani, R.T., Chowdri, I.N., and Dar, H. (2020). Factors influencing delay in initiating post-exposure prophylaxis for rabies prevention among animal bite victims: a cross sectional study. *J. Fam. Med. Prim. Care* 9, 4751–4755. https://doi.org/10.4103/jfmpc.jfmpc_890_20.
10. WHO; Rabies; Modelling Consortium (2019). The potential effect of improved provision of rabies post-exposure prophylaxis in Gavi-eligible countries: a modelling study. *Lancet Infect. Dis.* 19, 102–111. [https://doi.org/10.1016/S1473-3099\(18\)30512-7](https://doi.org/10.1016/S1473-3099(18)30512-7).
11. Hellgren, F., Cagigi, A., Arcoverde Cerveira, R., Ols, S., Kern, T., Lin, A., Eriksson, B., Dodds, M.G., Jasny, E., Schwendt, K., et al. (2023). Unmodified rabies mRNA vaccine elicits high cross-neutralizing antibody titers and diverse B cell memory responses. *Nat. Commun.* 14, 3713. <https://doi.org/10.1038/s41467-023-39421-5>.
12. Ng, W.M., Fedosyuk, S., English, S., Augusto, G., Berg, A., Thorley, L., Haselton, A.-S., Segireddy, R.R., Bowden, T.A., and Douglas, A.D. (2022). Structure of trimeric pre-fusion rabies virus glycoprotein in complex with two protective antibodies. *Cell Host Microbe* 30, 1219–1230.e7. <https://doi.org/10.1016/j.chom.2022.07.014>.
13. Cao, H., Li, H., Luan, N., Zhang, H., Lin, K., Hu, J., Song, J., and Liu, C. (2024). A rabies mRNA vaccine with H270P mutation in its glycoprotein induces strong cellular and humoral immunity. *Vaccine* 42, 1116–1121. <https://doi.org/10.1016/j.vaccine.2024.01.057>.
14. Fooks, A.R., Banyard, A.C., and Ertl, H.C.J. (2019). New human rabies vaccines in the pipeline. *Vaccine* 37, A140–A145. <https://doi.org/10.1016/j.vaccine.2018.08.039>.
15. Natesan, K., Isloor, S., Vinayagamurthy, B., Ramakrishnaiah, S., Doddamane, R., and Fooks, A.R. (2023). Developments in Rabies Vaccines: The Path Traversed from Pasteur to the Modern Era of Immunization. *Vaccines* 11, 756. <https://doi.org/10.3390/vaccines11040756>.
16. Sloan, S.E., Hanlon, C., Weldon, W., Niezgodna, M., Blanton, J., Self, J., Rowley, K.J., Mandell, R.B., Babcock, G.J., Thomas, W.D., et al. (2007). Identification and characterization of a human monoclonal antibody that potently neutralizes a broad panel of rabies virus isolates. *Vaccine* 25, 2800–2810. <https://doi.org/10.1016/j.vaccine.2006.12.031>.
17. Zorzan, M., Castellan, M., Gasparotto, M., Dias De Melo, G., Zecchin, B., Leopardi, S., Chen, A., Rosato, A., Angelini, A., Bourhy, H., et al. (2023). Antiviral mechanisms of two broad-spectrum monoclonal antibodies for

- rabies prophylaxis and therapy. *Front. Immunol.* *14*, 1186063. <https://doi.org/10.3389/fimmu.2023.1186063>.
18. Chao, T.-Y., Ren, S., Shen, E., Moore, S., Zhang, S.-F., Chen, L., Rupprecht, C.E., and Tsao, E. (2017). SYN023, a novel humanized monoclonal antibody cocktail, for post-exposure prophylaxis of rabies. *PLoS Negl. Trop. Dis.* *11*, e0006133. <https://doi.org/10.1371/journal.pntd.0006133>.
 19. de Melo, G.D., Hellert, J., Gupta, R., Corti, D., and Bourhy, H. (2022). Monoclonal antibodies against rabies: current uses in prophylaxis and in therapy. *Curr. Opin. Virol.* *53*, 101204. <https://doi.org/10.1016/j.coviro.2022.101204>.
 20. Thoulouze, M.I., Lafage, M., Schachner, M., Hartmann, U., Cremer, H., and Lafon, M. (1998). The Neural Cell Adhesion Molecule Is a Receptor for Rabies Virus. *J. Virol.* *72*, 7181–7190. <https://doi.org/10.1128/JVI.72.9.7181-7190.1998>.
 21. Khalifa, M.E., Unterholzner, L., and Munir, M. (2021). Structural and Evolutionary Insights Into the Binding of Host Receptors by the Rabies Virus Glycoprotein. *Front. Cell. Infect. Microbiol.* *11*, 736114. <https://doi.org/10.3389/fcimb.2021.736114>.
 22. Rustici, M., Bracci, L., Lozzi, L., Neri, P., Santucci, A., Soldani, P., Spreafico, A., and Niccolai, N. (1993). A model of the rabies virus glycoprotein active site. *Biopolymers* *33*, 961–969. <https://doi.org/10.1002/bip.360330612>.
 23. Gaudin, Y. (2000). Rabies Virus-Induced Membrane Fusion Pathway. *J. Cell Biol.* *150*, 601–612. <https://doi.org/10.1083/jcb.150.3.601>.
 24. Gaudin, Y., Tuffereau, C., Segretain, D., Knossow, M., and Flamand, A. (1991). Reversible conformational changes and fusion activity of rabies virus glycoprotein. *J. Virol.* *65*, 4853–4859. <https://doi.org/10.1128/JVI.65.9.4853-4859.1991>.
 25. Callaway, H.M., Zyla, D., Larrous, F., de Melo, G.D., Hastie, K.M., Avalos, R.D., Agarwal, A., Corti, D., Bourhy, H., and Saphire, E.O. (2022). Structure of the rabies virus glycoprotein trimer bound to a prefusion-specific neutralizing antibody. *Sci. Adv.* *8*, eabp9151. <https://doi.org/10.1126/sciadv.abp9151>.
 26. Yang, F., Lin, S., Ye, F., Yang, J., Qi, J., Chen, Z., Lin, X., Wang, J., Yue, D., Cheng, Y., et al. (2020). Structural Analysis of Rabies Virus Glycoprotein Reveals pH-Dependent Conformational Changes and Interactions with a Neutralizing Antibody. *Cell Host Microbe* *27*, 441–453.e7. <https://doi.org/10.1016/j.chom.2019.12.012>.
 27. Gaudin, Y., Ruigrok, R.W., Knossow, M., and Flamand, A. (1993). Low-pH conformational changes of rabies virus glycoprotein and their role in membrane fusion. *J. Virol.* *67*, 1365–1372. <https://doi.org/10.1128/JVI.67.3.1365-1372.1993>.
 28. Zhou, J., and Blissard, G.W. (2006). Mapping the conformational epitope of a neutralizing antibody (AcV1) directed against the AcMNPV GP64 protein. *Virology* *352*, 427–437. <https://doi.org/10.1016/j.virol.2006.04.041>.
 29. Siekavizza-Robles, C.R., Dollery, S.J., and Nicola, A.V. (2010). Reversible conformational change in herpes simplex virus glycoprotein B with fusion-from-without activity is triggered by mildly acidic pH. *Virol. J.* *7*, 352. <https://doi.org/10.1186/1743-422X-7-352>.
 30. Yao, Y., Ghosh, K., Epand, R.F., Epand, R.M., and Ghosh, H.P. (2003). Membrane fusion activity of vesicular stomatitis virus glycoprotein G is induced by low pH but not by heat or denaturant. *Virology* *310*, 319–332. [https://doi.org/10.1016/S0042-6822\(03\)00146-6](https://doi.org/10.1016/S0042-6822(03)00146-6).
 31. Roche, S., and Gaudin, Y. (2002). Characterization of the Equilibrium between the Native and Fusion-Inactive Conformation of Rabies Virus Glycoprotein Indicates That the Fusion Complex Is Made of Several Trimers. *Virology* *297*, 128–135. <https://doi.org/10.1006/viro.2002.1429>.
 32. Fan, L., Zhang, L., Li, J., and Zhu, F. (2022). Advances in the progress of monoclonal antibodies for rabies. *Hum. Vaccin. Immunother.* *18*, 2026713. <https://doi.org/10.1080/21645515.2022.2026713>.
 33. Ejemel, M., Smith, T.G., Greenberg, L., Carson, W.C., Lowe, D., Yang, Y., Jackson, F.R., Morgan, C.N., Martin, B.E., Kling, C., et al. (2022). A cocktail of human monoclonal antibodies broadly neutralizes North American rabies virus variants as a promising candidate for rabies post-exposure prophylaxis. *Sci. Rep.* *12*, 9403. <https://doi.org/10.1038/s41598-022-13527-0>.
 34. De Benedictis, P., Minola, A., Rota Nodari, E., Aiello, R., Zecchin, B., Salomoni, A., Foglierini, M., Agatic, G., Vanzetta, F., Lavenir, R., et al. (2016). Development of broad-spectrum human monoclonal antibodies for rabies post-exposure prophylaxis. *EMBO Mol. Med.* *8*, 407–421. <https://doi.org/10.15252/emmm.201505986>.
 35. Wang, W., Ma, J., Nie, J., Li, J., Cao, S., Wang, L., Yu, C., Huang, W., Li, Y., Yu, Y., et al. (2019). Antigenic variations of recent street rabies virus. *Emerg. Microbes Infect.* *8*, 1584–1592. <https://doi.org/10.1080/22221751.2019.1683436>.
 36. Dadonaite, B., Crawford, K.H.D., Radford, C.E., Farrell, A.G., Yu, T.C., Hannon, W.W., Zhou, P., Andrabi, R., Burton, D.R., Liu, L., et al. (2023). A pseudovirus system enables deep mutational scanning of the full SARS-CoV-2 spike. *Cell* *186*, 1263–1278.e20. <https://doi.org/10.1016/j.cell.2023.02.001>.
 37. Radford, C.E., Schommers, P., Gieselmann, L., Crawford, K.H.D., Dadonaite, B., Yu, T.C., Dingens, A.S., Overbaugh, J., Klein, F., and Bloom, J.D. (2023). Mapping the neutralizing specificity of human anti-HIV serum by deep mutational scanning. *Cell Host Microbe* *31*, 1200–1215.e9. <https://doi.org/10.1016/j.chom.2023.05.025>.
 38. Chun, T.W., Carruth, L., Finzi, D., Shen, X., DiGiuseppe, J.A., Taylor, H., Hermankova, M., Chadwick, K., Margolick, J., Quinn, T.C., et al. (1997). Quantification of latent tissue reservoirs and total body viral load in HIV-1 infection. *Nature* *387*, 183–188. <https://doi.org/10.1038/387183a0>.
 39. Pang, S., Koyanagi, Y., Miles, S., Wiley, C., Vinters, H.V., and Chen, I.S.Y. (1990). High levels of unintegrated HIV-1 DNA in brain tissue of AIDS dementia patients. *Nature* *343*, 85–89. <https://doi.org/10.1038/343085a0>.
 40. Sharkey, M.E., Teo, I., Greenough, T., Sharova, N., Luzuriaga, K., Sullivan, J.L., Bucy, R.P., Kostrikis, L.G., Haase, A., Veryard, C., et al. (2000). Persistence of episomal HIV-1 infection intermediates in patients on highly active anti-retroviral therapy. *Nat. Med.* *6*, 76–81. <https://doi.org/10.1038/71569>.
 41. Milone, M.C., and O'Doherty, U. (2018). Clinical use of lentiviral vectors. *Leukemia* *32*, 1529–1541. <https://doi.org/10.1038/s41375-018-0106-0>.
 42. Dong, W., and Kantor, B. (2021). Lentiviral Vectors for Delivery of Gene-Editing Systems Based on CRISPR/Cas: Current State and Perspectives. *Viruses* *13*, 1288. <https://doi.org/10.3390/v13071288>.
 43. Hislop, J.N., Islam, T.A., Eleftheriadou, I., Carpentier, D.C.J., Trabalza, A., Parkinson, M., Schiavo, G., and Mazarakis, N.D. (2014). Rabies Virus Envelope Glycoprotein Targets Lentiviral Vectors to the Axonal Retrograde Pathway in Motor Neurons. *J. Biol. Chem.* *289*, 16148–16163. <https://doi.org/10.1074/jbc.M114.549980>.
 44. Kondo, A., and Osawa, T. (2017). Establishment of an Extracellular Acidic pH Culture System. *J. Vis. Exp.* *129*, 56660. <https://doi.org/10.3791/56660>.
 45. Otwinowski, J., McCandlish, D.M., and Plotkin, J.B. (2018). Inferring the shape of global epistasis. *Proc. Natl. Acad. Sci. USA* *115*, E7550–E7558. <https://doi.org/10.1073/pnas.1804015115>.
 46. Beilstein, F., Abou Hamdan, A., Raux, H., Belot, L., Ouldali, M., Albertini, A.A., and Gaudin, Y. (2020). Identification of a pH-Sensitive Switch in VSV-G and a Crystal Structure of the G Pre-fusion State Highlight the VSV-G Structural Transition Pathway. *Cell Rep.* *32*, 108042. <https://doi.org/10.1016/j.celrep.2020.108042>.
 47. Hsieh, C.-L., Goldsmith, J.A., Schaub, J.M., DiVenere, A.M., Kuo, H.-C., Javanmardi, K., Le, K.C., Wrapp, D., Lee, A.G., Liu, Y., et al. (2020). Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science* *369*, 1501–1505. <https://doi.org/10.1126/science.abd0826>.
 48. Pallesen, J.L., Wang, N., Corbett, K.S., Wrapp, D., Kirchdoerfer, R.N., Turner, H.L., Cottrell, C.A., Becker, M.M., Wang, L., Shi, W., et al. (2017). Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc. Natl. Acad. Sci. USA* *114*, E7348–E7357. <https://doi.org/10.1073/pnas.1707304114>.

49. Wang, S., Li, R., Pan, X., Wang, M., Wu, Y., Li, Y., Huang, X., Zhu, R., Wang, X., Zhang, Y., et al. (2025). Structure-guided design of a prefusion GPC trimer induces neutralizing responses against LASV. *npj Vaccines* 10, 37. <https://doi.org/10.1038/s41541-025-01090-4>.
50. Ferlin, A., Raux, H., Baquero, E., Lepault, J., and Gaudin, Y. (2014). Characterization of pH-Sensitive Molecular Switches That Trigger the Structural Transition of Vesicular Stomatitis Virus Glycoprotein from the Postfusion State toward the Prefusion State. *J. Virol.* 88, 13396–13409. <https://doi.org/10.1128/JVI.01962-14>.
51. Yu, T.C., Thornton, Z.T., Hannon, W.W., DeWitt, W.S., Radford, C.E., Matsen, F.A., and Bloom, J.D. (2022). A biophysical model of viral escape from polyclonal antibodies. *Virus Evol.* 8, veac110. <https://doi.org/10.1093/ve/veac110>.
52. Nagarajan, T., Marissen, W., and Rupprecht, C. (2014). Monoclonal antibodies for the prevention of rabies: theory and clinical practice. *Antib. Technol. J.* 4, 1–12. <https://doi.org/10.2147/ANTI.S33533>.
53. Zhu, H., Chen, X., Shao, X., Ba, H., Wang, F., Wang, H., Yang, Y., Sun, N., Ren, J., Cheng, S., et al. (2015). Characterization of a virulent dog-originated rabies virus affecting more than twenty fallow deer (*Dama dama*) in Inner Mongolia, China. *Infect. Genet. Evol.* 31, 127–134. <https://doi.org/10.1016/j.meegid.2014.12.024>.
54. Franka, R., Carson, W.C., Ellison, J.A., Taylor, S.T., Smith, T.G., Kuzmina, N.A., Kuzmin, I.V., Marissen, W.E., and Rupprecht, C.E. (2017). In Vivo Efficacy of a Cocktail of Human Monoclonal Antibodies (CL184) Against Diverse North American Bat Rabies Virus Variants. *Trop. Med. Infect. Dis.* 2, 48. <https://doi.org/10.3390/tropicalmed2030048>.
55. Bakker, A.B.H., Python, C., Kissling, C.J., Pandya, P., Marissen, W.E., Brink, M.F., Lagerwerf, F., Worst, S., Van Corven, E., Kostense, S., et al. (2008). First administration to humans of a monoclonal antibody cocktail against rabies virus: Safety, tolerability, and neutralizing activity. *Vaccine* 26, 5922–5927. <https://doi.org/10.1016/j.vaccine.2008.08.050>.
56. Hellert, J., Buchrieser, J., Larrous, F., Minola, A., de Melo, G.D., Soriaga, L., England, P., Haouz, A., Telenti, A., Schwartz, O., et al. (2020). Structure of the prefusion-locking broadly neutralizing antibody RVC20 bound to the rabies virus glycoprotein. *Nat. Commun.* 11, 596. <https://doi.org/10.1038/s41467-020-14398-7>.
57. Gogtay, N.J., Munshi, R., Ashwath Narayana, D.H., Mahendra, B.J., Kshirsagar, V., Gunale, B., Moore, S., Cheslock, P., Thaker, S., Deshpande, S., et al. (2018). Comparison of a Novel Human Rabies Monoclonal Antibody to Human Rabies Immunoglobulin for Postexposure Prophylaxis: A Phase 2/3, Randomized, Single-Blind, Noninferiority, Controlled Study. *Clin. Infect. Dis.* 66, 387–395. <https://doi.org/10.1093/cid/cix791>.
58. Quiambao, B.P., Payumo, R.A., Roa, C., Borja-Tabora, C.F., Emmeline Montellano, M., Reyes, M.R.L., Zoleta-De Jesus, L., Capeding, M.R., Solimen, D.P., Barez, M.Y., et al. (2024). A phase 2b, Randomized, double blinded comparison of the safety and efficacy of the monoclonal antibody mixture SYN023 and human rabies immune globulin in patients exposed to rabies. *Vaccine* 42, 126018. <https://doi.org/10.1016/j.vaccine.2024.05.066>.
59. McClain, J.B., Chuang, A., Reid, C., Moore, S.M., and Tsao, E. (2021). Rabies virus neutralizing activity, pharmacokinetics, and safety of the monoclonal antibody mixture SYN023 in combination with rabies vaccination: Results of a phase 2, randomized, blinded, controlled trial. *Vaccine* 39, 5822–5830. <https://doi.org/10.1016/j.vaccine.2021.08.066>.
60. Synmore Biologics (Suzhou) Co., Ltd. (2023). A Phase III Randomized Blinded Study to Evaluate SYN023 Compared to Human Rabies Immune Globulin in Post Exposure Prophylaxis of Rabies in Adults With Category III Rabies Exposure Risks (clinicaltrials.gov). https://cdn.clinicaltrials.gov/large-docs/84/NCT04644484/Prot_000.pdf.
61. Goudsmit, J., Marissen, W.E., Weldon, W.C., Niezgodka, M., Hanlon, C.A., Rice, A.B., Kruijff, J. de Dietzschold, B., Bakker, A.B.H., and Rupprecht, C.E. (2006). Comparison of an Anti-Rabies Human Monoclonal Antibody Combination with Human Polyclonal Anti-Rabies Immune Globulin. *J. Infect. Dis.* 193, 796–801. <https://doi.org/10.1086/500470>.
62. Bakker, A.B.H., Marissen, W.E., Kramer, R.A., Rice, A.B., Weldon, W.C., Niezgodka, M., Hanlon, C.A., Thijsse, S., Backus, H.H.J., de Kruijff, J., et al. (2005). Novel human monoclonal antibody combination effectively neutralizing natural rabies virus variants and individual in vitro escape mutants. *J. Virol.* 79, 9062–9068. <https://doi.org/10.1128/JVI.79.14.9062-9068.2005>.
63. de Melo, G.D., Sonthonnax, F., Lepousez, G., Jouvion, G., Minola, A., Zatta, F., Larrous, F., Kergoat, L., Mazo, C., Moigneu, C., et al. (2020). A combination of two human monoclonal antibodies cures symptomatic rabies. *EMBO Mol. Med.* 12, e12628. <https://doi.org/10.15252/emmm.202012628>.
64. Marissen, W.E., Kramer, R.A., Rice, A., Weldon, W.C., Niezgodka, M., Faber, M., Slootstra, J.W., Meloen, R.H., Clijsters-van Der Horst, M., Visser, T.J., et al. (2005). Novel Rabies Virus-Neutralizing Epitope Recognized by Human Monoclonal Antibody: Fine Mapping and Escape Mutant Analysis. *J. Virol.* 79, 4672–4678. <https://doi.org/10.1128/JVI.79.8.4672-4678.2005>.
65. Kedari, A., Iheozor-Ejiofor, R., Salminen, P., Uğurlu, H., Mäkelä, A.R., Levanov, L., Vapalahti, O., Hytönen, V.P., Saksela, K., and Rissanen, I. (2024). Structural insight into rabies virus neutralization revealed by an engineered antibody scaffold. *Structure* 32, 2220–2230.e4. <https://doi.org/10.1016/j.str.2024.10.002>.
66. Dall'Acqua, W., Goldman, E.R., Lin, W., Teng, C., Tsuchiya, D., Li, H., Ysem, X., Braden, B.C., Li, Y., Smith-Gill, S.J., et al. (1998). A Mutational Analysis of Binding Interactions in an Antigen–Antibody Protein–Protein Complex. *Biochemistry* 37, 7981–7991. <https://doi.org/10.1021/bi980148j>.
67. Doud, M.B., Hensley, S.E., and Bloom, J.D. (2017). Complete mapping of viral escape from neutralizing antibodies. *PLoS Pathog.* 13, e1006271. <https://doi.org/10.1371/journal.ppat.1006271>.
68. Jin, L., Fendly, B.M., and Wells, J.A. (1992). High resolution functional analysis of antibody-antigen interactions. *J. Mol. Biol.* 226, 851–865. [https://doi.org/10.1016/0022-2836\(92\)90636-X](https://doi.org/10.1016/0022-2836(92)90636-X).
69. Pons, J., Rajpal, A., and Kirsch, J.F. (1999). Energetic analysis of an antigen/antibody interface: Alanine scanning mutagenesis and double mutant cycles on the hyHEL-10/lysozyme interaction. *Protein Sci.* 8, 958–968. <https://doi.org/10.1110/ps.8.5.958>.
70. Wang, Y., Rowley, K.J., Booth, B.J., Sloan, S.E., Ambrosino, D.M., and Babcock, G.J. (2011). G glycoprotein amino acid residues required for human monoclonal antibody RAB1 neutralization are conserved in rabies virus street isolates. *Antiviral Res.* 91, 187–194. <https://doi.org/10.1016/j.antiviral.2011.06.002>.
71. Kuzmina, N.A., Kuzmin, I.V., Ellison, J.A., and Rupprecht, C.E. (2013). Conservation of Binding Epitopes for Monoclonal Antibodies on the Rabies Virus Glycoprotein. *J. Antivir. Antiretrovir.* 5, 37–43. <https://doi.org/10.4172/jaa.1000061>.
72. Lentz, T.L. (1990). Rabies virus binding to an acetylcholine receptor α -subunit peptide. *J. Mol. Recognit.* 3, 82–88. <https://doi.org/10.1002/jmr.300030205>.
73. Ito, N., Okamoto, T., Sasaki, M., Miyamoto, S., Takahashi, T., Izumi, F., Inukai, M., Jarusombuti, S., Okada, K., Nakagawa, K., et al. (2021). Safety enhancement of a genetically modified live rabies vaccine strain by introducing an attenuating Leu residue at position 333 in the glycoprotein. *Vaccine* 39, 3777–3784. <https://doi.org/10.1016/j.vaccine.2021.05.002>.
74. Le Blois, H., Tuffereau, C., Blancou, J., Artois, M., Aubert, A., and Flamand, A. (1990). Oral immunization of foxes with avirulent rabies virus mutants. *Vet. Microbiol.* 23, 259–266. [https://doi.org/10.1016/0378-1135\(90\)90156-P](https://doi.org/10.1016/0378-1135(90)90156-P).
75. Nakagawa, K., Ito, N., Masatani, T., Abe, M., Yamaoka, S., Ito, Y., Okadera, K., and Sugiyama, M. (2012). Generation of a live rabies vaccine strain attenuated by multiple mutations and evaluation of its safety

- and efficacy. *Vaccine* 30, 3610–3617. <https://doi.org/10.1016/j.vaccine.2012.03.044>.
76. Itakura, Y., Tabata, K., Morimoto, K., Ito, N., Chambaro, H.M., Eguchi, R., Otsuguro, K.I., Hall, W.W., Orba, Y., Sawa, H., et al. (2022). Glu333 in rabies virus glycoprotein is involved in virus attenuation through astrocyte infection and interferon responses. *iScience* 25, 104122. <https://doi.org/10.1016/j.isci.2022.104122>.
 77. Tao, L., Ge, J., Wang, X., Zhai, H., Hua, T., Zhao, B., Kong, D., Yang, C., Chen, H., and Bu, Z. (2010). Molecular basis of neurovirulence of flury rabies virus vaccine strains: importance of the polymerase and the glycoprotein R333Q mutation. *J. Virol.* 84, 8926–8936. <https://doi.org/10.1128/JVI.00787-10>.
 78. Hadfield, J., Megill, C., Bell, S.M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P., Bedford, T., and Neher, R.A. (2018). Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* 34, 4121–4123. <https://doi.org/10.1093/bioinformatics/bty407>.
 79. Pattnaik, P., Mahal, A., Mishra, S., Alkhoury, A., Mohapatra, R.K., and Kandi, V. (2023). Alarming Rise in Global Rabies Cases Calls for Urgent Attention: Current Vaccination Status and Suggested Key Countermeasures. *Cureus* 15, e50424. <https://doi.org/10.7759/cureus.50424>.
 80. Haddock, H.K., Dingens, A.S., Hilton, S.K., Overbaugh, J., and Bloom, J.D. (2018). Mapping mutational effects along the evolutionary landscape of HIV envelope. *eLife* 7, e34420. <https://doi.org/10.7554/eLife.34420>.
 81. Starr, T.N., Greaney, A.J., Hannon, W.W., Loes, A.N., Hauser, K., Dillen, J.R., Ferri, E., Farrell, A.G., Dadonaite, B., McCallum, M., et al. (2022). Shifting mutational constraints in the SARS-CoV-2 receptor-binding domain during viral evolution. *Science* 377, 420–424. <https://doi.org/10.1126/science.abo7896>.
 82. Lei, R., Liang, W., Ouyang, W.O., Hernandez Garcia, A., Kikuchi, C., Wang, S., McBride, R., Tan, T.J.C., Sun, Y., Chen, C., et al. (2024). Epistasis mediates the evolution of the receptor binding mode in recent human H3N2 hemagglutinin. *Nat. Commun.* 15, 5175. <https://doi.org/10.1038/s41467-024-49487-4>.
 83. Bonnaud, E.M., Troupin, C., Dacheux, L., Holmes, E.C., Monchatre-Leroy, E., Tanguy, M., Bouchier, C., Cliquet, F., Barrat, J., and Bourhy, H. (2019). Comparison of intra- and inter-host genetic diversity in rabies virus during experimental cross-species transmission. *PLoS Pathog.* 15, e1007799. <https://doi.org/10.1371/journal.ppat.1007799>.
 84. Schnell, M.J., McGettigan, J.P., Wirblich, C., and Papaneri, A. (2010). The cell biology of rabies virus: using stealth to reach the brain. *Nat. Rev. Microbiol.* 8, 51–61. <https://doi.org/10.1038/nrmicro2260>.
 85. Gluska, S., Zahavi, E.E., Chein, M., Gradus, T., Bauer, A., Finke, S., and Perlson, E. (2014). Rabies Virus Hijacks and Accelerates the p75NTR Retrograde Axonal Transport Machinery. *PLoS Pathog.* 10, e1004348. <https://doi.org/10.1371/journal.ppat.1004348>.
 86. Tordo, N., and Poch, O. (1988). Structure of Rabies Virus. In *Rabies Developments in Veterinary Virology*, J.B. Campbell and K.M. Charlton, eds. (Springer), pp. 25–45. https://doi.org/10.1007/978-1-4613-1755-5_2.
 87. Nitschel, S., Zaack, L.M., Potratz, M., Nolden, T., Te Kamp, V., Franzke, K., Höper, D., Pfaff, F., and Finke, S. (2021). Point Mutations in the Glycoprotein Ectodomain of Field Rabies Viruses Mediate Cell Culture Adaptation through Improved Virus Release in a Host Cell Dependent and Independent Manner. *Viruses* 13, 1989. <https://doi.org/10.3390/v13101989>.
 88. Yamada, K., Noguchi, K., Nonaka, D., Morita, M., Yasuda, A., Kawazato, H., and Nishizono, A. (2013). Addition of a single N-glycan to street rabies virus glycoprotein enhances virus production. *J. Gen. Virol.* 94, 270–275. <https://doi.org/10.1099/vir.0.047852-0>.
 89. Hannon, W.W., and Bloom, J.D. (2024). dms-viz: Structure-informed visualizations for deep mutational scanning and other mutation-based datasets. *J. Open Source Softw.* 9, 6129. <https://doi.org/10.21105/joss.06129>.
 90. Loes, A.N., Tarabi, R.A.L., Huddleston, J., Touyon, L., Wong, S.S., Cheng, S.M.S., Leung, N.H.L., Hannon, W.W., Bedford, T., Cobey, S., et al. (2024). High-throughput sequencing-based neutralization assay reveals how repeated vaccinations impact titers to recent human H1N1 influenza strains. *J. Virol.* 98, e0068924. <https://doi.org/10.1128/jvi.00689-24>.
 91. Crawford, K.H.D., and Bloom, J.D. (2019). alignparse: A Python package for parsing complex features from high-throughput long-read sequencing. *J. Open Source Softw.* 4, 1915. <https://doi.org/10.21105/joss.01915>.
 92. Haddock, H.K., Galloway, J.G., Dadonaite, B., Bloom, J.D., Matsen, F.A., and DeWitt, W.S. (2023). Jointly modeling deep mutational scans identifies shifted mutational effects among SARS-CoV-2 spike homologs. Preprint at bioRxiv. <https://doi.org/10.1101/2023.07.31.551037>.
 93. Corti, D. (2016). Antibodies That Potently Neutralize Rabies Virus and Other Lyssaviruses and Uses Thereof. WIPO patent WO2016078761A1.
 94. Bakker, A.B.H., and Marissen, W.E. (2011). Liquid Anti-rabies Antibody Formulations. European patent EP2088997B1.
 95. Shen, E., and Ren, S. (2013). Compositions and Methods Related to Prevention and Treatment of Rabies Infection. WIPO patent WO2013174003A1.
 96. Larsen, B.B., McMahon, T., Brown, J.T., Wang, Z., Radford, C.E., Crowe, J.E., Velesler, D., and Bloom, J.D. (2024). Functional and antigenic landscape of the Nipah virus receptor binding protein. Preprint at bioRxiv. <https://doi.org/10.1101/2024.04.17.589977>.
 97. Carr, C.R., Crawford, K.H.D., Murphy, M., Galloway, J.G., Haddock, H.K., Matsen, F.A., Andersen, K.G., King, N.P., and Bloom, J.D. (2024). Deep mutational scanning reveals functional constraints and antibody-escape potential of Lassa virus glycoprotein complex. *Immunity* 57, 2061–2076. e11. <https://doi.org/10.1016/j.immuni.2024.06.013>.
 98. Dadonaite, B., Brown, J., McMahon, T.E., Farrell, A.G., Figgins, M.D., Asarnow, D., Stewart, C., Lee, J., Logue, J., Bedford, T., et al. (2024). Spike deep mutational scanning helps predict success of SARS-CoV-2 clades. *Nature* 631, 617–626. <https://doi.org/10.1038/s41586-024-07636-1>.
 99. Dadonaite, B., Ahn, J.J., Ort, J.T., Yu, J., Furey, C., Dosey, A., Hannon, W.W., Vincent Baker, A.L., Webby, R.J., King, N.P., et al. (2024). Deep mutational scanning of H5 hemagglutinin to inform influenza virus surveillance. *PLoS Biol.* 22, e3002916. <https://doi.org/10.1371/journal.pbio.3002916>.
 100. Humes, D., Rainwater, S., and Overbaugh, J. (2021). The TOP vector: a new high-titer lentiviral construct for delivery of sgRNAs and transgenes to primary T cells. *Mol. Ther. Methods Clin. Dev.* 20, 30–38. <https://doi.org/10.1016/j.omtm.2020.10.020>.
 101. Huddleston, J., Hadfield, J., Sibley, T.R., Lee, J., Fay, K., Ilcisin, M., Harkins, E., Bedford, T., Neher, R.A., and Hodcroft, E.B. (2021). Augur: a bioinformatics toolkit for phylogenetic analyses of human pathogens. *J. Open Source Softw.* 6, 2906. <https://doi.org/10.21105/joss.02906>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
17C7	Synthesized by Genscript	Patent US8226952B2
CR4098	Synthesized by Genscript	Patent US7959922B2
RVA122	Synthesized by Genscript	Patent WO2016/078761A1
RVC58	Synthesized by Genscript	Patent WO2016/078761A1
CR57	Synthesized by Genscript	Patent US7959922B2
RVC20	Synthesized by Genscript	See Hellert et al. ⁵⁶
RVC68	Synthesized by Genscript	Patent WO2016/078761A1
CTB012	Synthesized by Genscript	Patent WO2013174003A1
Goat anti-Human IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647	Invitrogen	Ref No. A-21445; RRID:AB_2535862
Bacterial and virus strains		
10-beta electrocompetent cells	New England Biolabs	Ref No. C3020K
Chemicals, peptides, and recombinant proteins		
Puromycin	Sigma	Ref No. P8833-25mg
Trypsin EDTA	Fisher	Ref No. MT25053CI
DMEM	Fisher	Ref No. MT10013CV
Tetracycline-free fetal bovine serum	Gemini Bio	Ref No. 100-800
Penicillin-Streptomycin	Fisher	Ref No. MT30002CI
L-glutamine	Fisher	Ref No. MT25005CI
BioT Transfection Reagent	Bioland Scientific	Ref No. B01-02
Doxycycline hyclate	Tocris Bioscience	Ref No. 40-905-0
DMEM powder	Corning	Ref No. 90013PB
Sodium bicarbonate	Sigma	Ref No. S5761-500G
Poly-L-Lysine solution	Sigma	Ref No. 4707-50ML
Polybrene	EMD Millipore	Ref No. TR1003-G
Cell Dissociation Buffer Enzyme-Free PBS-Based	Gibco	Ref No. 13151-014
Paraformaldehyde (4% in PBS)	Biotium	Ref No. 22023
Bovine Serum Albumin Fraction V	Gibco	Ref No. 15260-037
Critical commercial assays		
QIAprep Spin Miniprep Kit	Qiagen	Ref No. 27106X4
HiSpeed Plasmid Maxi Kit	Qiagen	Ref No. 12662
Bright-Glo Luciferase Assay System	Promega	Ref No. E2610
Qubit 1x dsDNA High Sensitivity and Broad Range Assay Kits	Fisher	Ref No. Q33231
NEBuilder HiFi DNA Assembly Kit	New England Biolabs	Ref No. E2621
KOD Hot Start Master Mix	ThermoFisher	Ref No. 7182-4
KLD Enzyme Mix	New England Biolabs	Ref No. M0554
Deposited data		
Data generated by vep-dms-pipeline-3	This paper	https://dms-vep.org/RABV_Pasteur_G_DMS/
Experimental models: Cell lines		
293T cells	ATCC	Ref No. CRL-3216
293T cells expressing rTA	Dadonaite et al. ³⁶	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Oligonucleotides		
Primers for plasmid cloning, mutagenesis, amplification of viral templates by PCR, and next-generation sequencing library preparation	IDT	Sequences at https://github.com/dms-vep/RABV_Pasteur_G_DMS/tree/main/Additional_Data/PrimerSequences
Recombinant DNA		
Expression plasmid for VSV-G (HDM_VSV_G)	Addgene	Plasmid #204156
Expression plasmid for Pasteur strain rabies G	This paper	Sequence at https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/3968_HDM_Rabies-G_Pasteur_GenscriptOpt3.gb
HDM-Hgpm2	Addgene	Plasmid #204152
pRC-CMV-Rev1b	Addgene	Plasmid #204153
HDM-tat1b	Addgene	Plasmid #204154
Lentiviral backbone for Pasteur strain rabies G	This paper	Sequence at https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/3969_V5LP_pH2rU3_ForInd_ExtGag_Humes_RABV-G_Pasteur-Genscript3_CMV_zsGT2APurR.gb
Expression plasmid for A12_2718 strain rabies G	This paper	Sequence at https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/4956_HDM_Rabies-G_AGN94399-WT.gb
Expression plasmid for NY-2011-12020 strain rabies G	This paper	Sequence at https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/4959_HDM_Rabies-G_QEJ74619_WT.gb
Lentiviral backbone for zsGreen and Luc2	Addgene	Plasmid #164432
Software and algorithms		
vep-dms-pipeline-3	N/A	https://github.com/dms-vep/dms-vep-pipeline-3
polyclonal	Yu et al. ⁵¹	https://github.com/jbloomlab/polyclonal
neutcurve	Loes et al. ⁹⁰	https://github.com/jbloomlab/neutcurve
alignparse	Crawford and Bloom ⁹¹	https://github.com/jbloomlab/alignparse
multidms	Haddox et al. ⁹²	https://github.com/matsengrp/multidms

EXPERIMENTAL MODEL DETAILS

Rabies G used for deep mutational scanning

We performed the deep mutational scanning using a codon-optimized G protein from the Pasteur strain (NCBI reference number: NC_001542.1, locus number: NP_056796), which is a vaccine strain¹⁵ with a G protein that has had its structure determined by cryo-EM.^{12,25} See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/LibrarySpecs/Rabies-G_Pasteur-fullORF.fasta for the full length codon-optimized G used in this study. To maintain consistency with the standard numbering convention for G,^{19,25,56} we assign the number 1 to the first protein site in the ectodomain (site 20 in the full length protein encoded by the G ORF). Note that the interactive Nextstrain trees at <https://nextstrain.org/groups/jbloomlab/dms/rabies-G> have options to number sites either using this standard G ectodomain numbering sequence or starting with the N-terminal methionine of the full-length protein.

Cells

As target cells for infections to measure mutation effects on cell entry and antibody escape, we used 293T cells from ATCC (CRL-3216). For producing pseudovirus libraries from integrated cells (Figure S1), we used a previously described³⁶ 293T-rtTA cell line. We cultured all cell lines in D10 media (Dulbecco's Modified Eagle Medium supplemented with 10% heat-inactivated fetal bovine serum, 2 mM l-glutamine, 100 U/mL penicillin, and 100 mg/mL streptomycin). When working with the 293T-rtTA cell line, we used tetracycline-negative fetal bovine serum (Gemini Bio, Ref. No. 100-800). For generating pseudovirus libraries that either express

rabies G mutants or VSV-G, we used Dulbecco's Modified Eagle Medium without phenol red as we found this was helpful for pseudovirus concentration as it avoided also concentrating phenol red.

We noted that titers for pseudovirus expressing rabies G were higher at a pH of 7.1 (Figure S2B). For all infections with the rabies G pseudovirus libraries and single mutant validation assays, we formulated cell culture media that adopts a pH of 7.1 at 37°C and 5% CO₂ based on a prior published protocol,⁴⁴ supplemented with 4 µg/mL polybrene (EMD Millipore, Ref. No. TR-1003-G). We used powdered Dulbecco's Modified Eagle Medium (Corning™ 90013PB) formulated without L-glutamine, pyruvate, sodium bicarbonate, and phenol red. This media base was reconstituted in Milli-Q water per manufacturer's instructions and supplemented with 10% tetracycline-negative heat-inactivated fetal bovine serum, 2 mM L-glutamine, 100 U/mL penicillin, 100 mg/mL streptomycin, and sodium bicarbonate to a final concentration of ~0.75 g/L. Prior to use, we confirmed the pH of media after incubation for 18 hours 37°C and 5% CO₂.

Antibodies

Antibodies were produced by Genscript as human IgG1 kappa isotypes. The sequences for heavy and light chains are at (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/AntibodySequences/RABV-G_DMS_AntibodySequences.fasta). Sequences were obtained from original patents or publications. RVA122, RVC58 and RVC58 originated from patent WO2016/078761A1,⁹³ CR4098 and CR57 originated from patent US7959922B2,⁹⁴ RVC20 came from original publication,⁵⁶ and CTB012 from patent WO2013174003A1.⁹⁵

METHOD DETAILS

G plasmid mutant libraries

The lentiviral backbone is schematized in Figure S1A and is mostly identical to the backbone previously used for pseudovirus deep mutational scanning of other viral entry proteins.^{36,37,96–99} However, to enhance packaging of genomes in pseudovirus particles, we modified the lentiviral backbone to contain an extended Gag sequence.¹⁰⁰ A full plasmid map that contains the lentiviral backbone encoding the codon-optimized rabies G sequence is at (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/3969_V5LP_pH2rU3_ForInd_ExtGag_Humes_RABV-G_Pasteur-Genscript3_CMV_zsGT2APurR.gb).

We mutagenized G starting two residues prior to the beginning of the ectodomain and continuing up to site 431 (so sites -2 to 431 in the standard numbering scheme). We designed the mutant library to encode all 19 mutant amino-acids at each site, and also to include 20 stop codons at alternating positions from the start of the mutagenized region as negative controls for cell entry. We ordered the mutant library as a site-saturation variant library from Twist Biosciences. The final Twist quality control report for the library is available at ([https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/LibrarySpecs/Q-293667_Final_QC_Report%20\(9\).xlsx](https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/LibrarySpecs/Q-293667_Final_QC_Report%20(9).xlsx)). We noted that 38 mutants from the overall library were missing. Only site H384 (site 403 in Quality Control report) failed in synthesis, with 14 amino acid mutations missing.

We appended barcodes consisting of 16 random nucleotides to the mutagenized rabies G gene fragments downstream of the stop codon using PCR largely as described previously.⁹⁶ To generate biological replicate libraries, Library A and Library B, we performed two separate barcoding PCR reactions. For each barcoding PCR, we combined 5 ng of the library template from Twist (1 µL), 1.5 µL of forward primer (BC_FOR, 10 µM stock), and 1.5 µL of reverse primer (BC_REV, 10 µM stock), 21 µL of molecular biology grade water with KOD Hot Start Master Mix (ThermoFisher, Ref. No. 71842-4). See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV-G_Sequencing-Primers.fasta primer sequences. PCR cycling conditions were as follows:

95°C, 2 min
95°C, 20 sec
55.5°C, 20 sec, cooling at 0.5°C/sec
70°C, 1 min
Return to Step 2, 9 cycles
12°C hold

The PCR product was run on a 0.8% agarose gel and a fragment of the correct size was excised for clean up using a NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel, Ref. No. 740609.5) followed by Ampure XP beads (Beckman Coulter, Ref. No. A63881) and elution in water.

To prepare the recipient lentiviral vector, we digested a lentiviral plasmid that contains mCherry in the location where the G viral entry protein is cloned (4016_V5LP_pHrU3_ForInd-Extgag_mcherry) with XbaI and MluI-HF (NEB, Cat. No. R0145S and R3198S respectively). The digested product was gel purified and cleaned with Ampure XP beads as for the PCR product.

For each library, barcoded mutants were cloned into the digested lentiviral vector using HiFi assembly at a 2:1 insert to vector molar ratio, at 50°C for 1 hour. HiFi products for each library were cleaned with Ampure XP beads and transformed into 10-beta electro-competent cells (NEB, Ref. No. C3020K) with a MicroPulser Electroporator (BioRad, Ref. No. 1652100), shocking at 2 kV for 5 milliseconds. Transformants were plated onto LB+ampicillin plates overnight at 37°C. To ensure the library was not bottlenecked and that the barcodes that are ultimately assigned to different variants are unique, we sought to maximize the number of colonies for each

library. (Note that the library size is determined at the step of creating the cell-integrated libraries as schematized in [Figure S1B](#), not at the plasmid creation stage; the plasmid library size must substantially exceed the final desired size of the cell-integrated libraries as there can be recombination during production of the lentiviruses that are integrated into the 293T-rtTA cells.) Library A contained 9.2×10^6 colonies, and Library B contained 7.8×10^6 colonies. Colonies were scraped off of plates into liquid luria broth, and plasmids were maxi-prepped with a HiSpeed Plasmid Maxi Kit (Qiagen, Ref. No. 12662).

Cell-stored rabies G mutant library

A deep mutational scan requires the ability to link the rabies G mutant expressed on each pseudovirus's surface with the barcode encoded in the genome. To accomplish this, we used the previously described^{36,37,96,97,99} approach shown in [Figure S1B](#), which involves creating a library of cells each containing a single integrated lentiviral genome encoding a different barcoded G mutant. As schematized in [Figure S1B](#), we first generated VSV-G expressing pseudoviruses encoding the rabies G mutants in their genomes. Using VSV-G as the viral glycoprotein ensures that cell entry occurs independently of the function of the rabies G encoded in the lentiviral genome. Next we used these VSV-G pseudovirions to transduce the 293T-rtTA cell line at a low multiplicity of infection (<0.01) to ensure single genome integration into these cells, followed by selection using puromycin (Sigma, Ref. No. P8833-25MG).

Specifically we produced VSV-G pseudovirus that package the rabies G encoding genomes by transfection at the scale of 30 mL. Approximately 4×10^6 cells were plated in each of three 10 cm dishes. Each plate was transfected with 5 μ g of the plasmid encoding the lentiviral backbone with barcoded G mutants, 1.25 μ g each of plasmids 26_HDM-Hgpm2 (Gagpol), 27_HDM-tat1b (Tat), 28_pRC-CMV-Rev1b (Rev), and 29_HDM-VSV-G (these plasmids are all available on AddGene under IDs 204152, 204154, 204153, and 204156; see <https://www.addgene.org/browse/article/28237987/>). Plasmids were transfected using the BioT transfection reagent (Bioland Scientific, Ref. No. B01-02) per manufacturer's instructions. After 48 hours, supernatant from the plates was collected and filtered using a 0.45 μ m syringe filter (Corning, Ref. No. 431220), and this supernatant was stored at -80°C until further use. To measure transducing units/mL (TU/mL) of this virus, we thawed an aliquot and transduced 293T cells and measured counts of ZsGreen-positive cells by flow cytometry.

As there were $\sim 8,000$ amino acid mutations in the rabies G library, we aimed for 80,000 variants so that each mutant would have approximately 10 barcodes. We followed a prior protocol³⁷ as described to transduce 293T-rtTA cells and collect transduced cells. Briefly, 6-well plates were seeded with 600,000 293T-rtTA cells. At the time of infection, we counted cells in a subset of wells and calculated transducing units of VSV-G pseudovirus to achieve a multiplicity of infection of 0.01. We transduced approximately 25 wells of cells for each library. After 48 hours, we measured transduction for some wells by flow cytometry counting ZsGreen cells and observed a final multiplicity of infection of 0.006. Using this percentage and the cell counts at the time of infection, we back-calculated the number of initial infected cells per well and pooled cells from wells to achieve approximately 80,000 infected cells for each library. Note that this entire process was performed separately for each of the two duplicate barcoded plasmid libraries described above.

Each library was pooled into 50 mL aliquots in tetracycline-negative D10 media supplemented with 0.75 μ g/mL puromycin to select for transduced cells, which constitutively express the puromycin resistance gene in the lentiviral genome. These cells were initially plated in two 15 cm plates for each library. Cells were passaged and expanded in tetracycline-negative media with 0.75 μ g/mL puromycin until only ZsGreen-positive cells were observed by microscopy. To expand the cell libraries, each was expanded into four 5-layer flasks (Falcon, Ref. No. 353144) in the absence of puromycin. To prevent future bottlenecks, we passaged cells at numbers in vast excess of the variants (eg. minimum 20×10^6 cells). Cell libraries were stored in aliquots of over 20×10^6 cells per aliquot in the vapor phase of liquid nitrogen. To protect the cell libraries during the freeze-thaw process, we froze cells in freezing media, composed of tetracycline-negative D10 media supplemented with 10% DMSO (by volume) and an additional 10% (by volume) of tetracycline-negative fetal bovine serum.

Production of rabies G and VSV-G expressing pseudovirus libraries

After creating the library of cells each of which encodes only one barcoded rabies G mutant lentiviral backbone, we produced pseudovirus particles expressing either rabies G or VSV-G on their surface ([Figure S1B](#)). The VSV-G expressing pseudoviruses were used for sequencing to link the barcode to the rabies G mutants (see below) and to control for variation in library composition during experiments to measure the effects of mutations on cell entry.

To produce rabies G the pseudovirus library, we seeded a minimum of three 5-layer flasks with ~ 150 million cells each in phenol-free, tetracycline-negative D10 media. To each flask, doxycycline was added to a final concentration of 1 μ g/mL to begin inducing expression of rabies G mutants from integrated genomes. At the time of transfecting helper plasmids 24 hours later, the D10 media was swapped for ~ 150 mL of fresh media supplemented with 1 μ g/mL of doxycycline. For each flask, 60 μ g each of 26_HDM-Hgpm2 (Gagpol), 27_HDM-tat1b (Tat), and 28_pRC-CMV-Rev1b (Rev) expression plasmids were combined with 9.2mL of DMEM and 276 μ L of BioT transfection reagent. This transfection mix was incubated at room temperature for 15 minutes then added to the 5-layer flasks. Approximately 40 hours later, media supernatant was collected and sterile filtered using 0.45 μ m SFCA 500mL Rapid-Flow filter units (Nalgene, Ref. No. 09-740-44B). Filtered supernatant was concentrated ~ 10 -fold using Pierce™ PES, 100K MWCO, 20–100 mL Protein Concentrators (Thermo, Ref. No. 88537) by centrifugation at $\sim 1210g$ for 30–40 minutes, which was sufficient to achieve pseudovirus library titers of 10^6 transduction units/mL. Until further use, the virus was aliquoted and stored at -80°C .

We produced VSV-G expressing pseudovirus by seeding two 15 cm plates each with 20 million library cells at the same time of seeding 5-layer flasks. To reduce expression of rabies G in these cells, we seeded cells in tetracycline-negative media and

maintained cells in tetracycline-negative media when producing pseudovirus (note some rabies G will still be expressed off the lentiviral Tat-driven promoter). As with rabies G library pseudovirus, we replaced media immediately prior to transfection. To each plate, we transfected 9 μ g each of HDM_Hgpm2 (Gagpol), 27_HDM_tat1b (Tat), and pRC_CMV_Rev1b (Rev) expression plasmids with 3.75 μ g of 29_HDM_VSV-G expression plasmids. These plasmids were combined with 1.5mL of DMEM and 45 μ L of BioT. After incubation for 15 minutes, this transfection was added drop-wise to cells as per manufacturer instructions. Forty hours later, supernatant was sterile filtered using a 0.45 μ m syringe filter (Corning, Ref. No. 431220) and aliquoted for storage at -80°C.

To determine the necessary volume of virus for experiments, we measured titers in transducing units/mL (TU/mL). We note that measuring titers may underestimate the viral particles used in all assays. Subsequent experiments rely on mini-prepping unintegrated lentiviral genomes, which are in excess of integrated genomes measured by titers.^{38–40} However, we utilized titration measurements to avoid the risk of bottlenecking the library during infection experiments.

For viral titer measurements, small aliquots of virus were thawed at room temperature. To determine the TU/ml, we made a 5-fold dilution series of virus and measured the percentage of 293T cells infected at multiple concentrations of virus using flow cytometry for the ZsGreen encoded in the lentiviral genome at 40–48 hours post-infection. For VSV-G library pseudovirus, we performed these infections in standard tetracycline-free D10 media (pH of 7.4). For rabies G pseudovirus, we diluted virus in pH 7.1 media and swapped cell media for pH 7.1 media because (as described above) low pH media yielded better titers for rabies G. Prior to combining rabies G library pseudovirus and cells, media was equilibrated to the correct pH at 37°C and 5% CO₂ in a tissue culture incubator for 1 hour. For antibody escape infection assays described later, we also measured barcoded VSV-G mCherry pseudovirus in these low pH conditions to plan the volume required to achieve 1–2% of titers relative to rabies G pseudovirus.

Long-read sequencing to link barcodes with rabies G mutants

We used PacBio long-read sequencing to link barcodes with the encoded rabies G variants in the lentiviral genome, largely as described previously.^{36,37,96,97} Because the pseudo-diploid nature of HIV-derived lentiviruses leads to recombination during the infection of 293T-rtTA cells for generating single integrants, we performed the PacBio sequencing on pseudoviruses generated from the integrated cells rather than the plasmids initially used to generate the pseudovirus to create these cells.³⁶

Specifically, 24 hours prior to infections, we plated ~500,000–600,000 293T cells per well in poly-L-lysine coated 6-well plates. We aimed to transduce cells with ~12 million transducing units of VSV-G expressing pseudovirus for each library. VSV-G expressing pseudovirus for each library was separately concentrated by centrifugation using Amicon Ultra 15mL/100kD MWCO Centrifugal filters (EMD Millipore, Ref. No. UFC9100024). These concentrated viruses were resuspended to a volume of ~7mL of D10 media. We removed media from 3 wells of cells and added 2mL of virus to each well. After infection for 12–14 hours, each well of cells was separately miniprep using a QIAprep Spin Miniprep Kit (Qiagen, Ref. No. 2710) to recover non-integrated reverse-transcribed lentiviral DNA. To improve yield, we heated miniprep columns during the elution step to 55°C for ~5 minutes prior to centrifugation. After elution, we pooled the miniprep eluent each library for subsequent PCR steps.

We performed a two-step PCR product to generate amplicons for long-read sequencing as originally described in Dadonaite et al.³⁶ To detect strand exchange events originating from the PCR, we performed a first round of PCR in two separate reactions that introduces a single nucleotide tag (either a G or C) at the 5' or 3' ends of the amplicon. For each reaction, we combined 20 μ L of KOD Hot Start Master Mix, 15 μ L of miniprep DNA, and molecular biology grade water. For primers, we either added 1 μ L each of 5_PacBio_primer_G and 3_PacBio_primer_C (both at 10 μ M stock) or 1 μ L each of 5_PacBio_primer_C and 3_PacBio_primer_G (both at 10 μ M stock). See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV_G_Sequencing-Primers.fasta for PacBio Round 1 primers. PCR cycling conditions were as follows:

95°C, 2 min
95°C, 20 sec
60°C, 10 s, cooling at 0.5°C/sec
70°C, 1 min
Return to Step 2, 7 cycles
70°C, 1 min
4°C hold

The PCR products were cleaned with 50 μ L of Ampure XP beads (1:1 volume ratio with PCR reaction) and eluted in 35 μ L of elution buffer. For each library, we pooled equal volumes of each separate PCR reaction into a template master mix for the second step of the PCR. For each step 2 PCR reaction, we combined 21 μ L of the round 1 product, with 2 μ L each of 5_PacBio_Rnd2 (10 μ M stock) and 3_PacBio_Rnd2 (10 μ M stock) primers. See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV_G_Sequencing-Primers.fasta for the PacBio Round 2 primer sequences. PCR cycling conditions were as follows:

95°C, 2 min
95°C, 20 sec
70°C, 1 sec
60°C, 10 s, cooling at 0.5°C/sec

70°C, 1 min
Return to Step 2, 10 cycles
70°C, 1 min
4°C hold

Each PCR reaction was cleaned with 50 μ L of Ampure XP beads (1:1 volume ratio with PCR reaction) and eluted in 40 μ L of elution buffer. We combined PCR reactions for each library together and verified by TapeStation that the amplicons were monodisperse and of correct length prior to sequencing. We sequenced each library on a single SMRTcell with a movie length of 30 hours on a PacBio Sequel IIe sequencer. To sequence to a higher depth, we later sequenced Library B on an additional SMRTcell and appended this to our analysis. Computational analysis of long-read sequencing to establish the barcode and variant linkages is detailed in the “[PacBio sequencing analysis](#)” section. That analysis checked for swapping of the tags added in the first PCR on the same molecule to detect strand exchange; as described in the data analysis, the rate of strand exchange was low presumably because we carefully limited the number of PCR cycles.

Deep mutational scanning for mutational effects on cell entry

To measure effects of G mutations on cell entry in 293T cells, we infected 293T cells with the rabies G pseudovirus library. To correct for library composition bias, we simultaneously infected a separate set of 293T cells with VSV-G expressing library pseudovirus ([Figure S2A](#)). Because VSV-G drives infection in the latter case, all pseudoviruses regardless of rabies G mutant function can mediate cell entry when VSV-G is present.

For all infection assays, we coated 6-well plates (Corning, Ref. No. 3516) with poly-L-lysine (Sigma-Aldrich, Ref. No. P4707-50mL). To do this, we added 1mL of poly-L-lysine to all plate wells and ensured full coating. After a 5 minute incubation, poly-L-lysine was aspirated, wells were washed with molecular biology grade water and allowed to dry at room temperature. We seeded each well with between 700,000 to 800,000 293T cells. Approximately 24 hours later after seeding cells, infection assays were performed.

For the rabies G pseudovirus library, we wanted each replicate well for the infection assays to have over 10-fold variant coverage in our library to prevent bottlenecks, so we infected with >1 million transducing units of virus per replicate well. (Note that the actual number of sequenced variants that enter cells may be higher, as transduction units are determined by flow cytometry which measured integrated viral genomes, but the non-integrated lentiviral DNA we mini-prepare for sequencing is generally in substantial excess to integrated viral genomes.^{38–40}) For our infection assays, rabies G pseudovirus library was first exchanged into D10 media adjusted to a pH of 7.1 and supplemented with 4 μ g/mL polybrene. To do this, we added 2 times the volume of pH 7.1 media to the virus used in a particular experiment. This was then spin concentrated at 3100g for 30 minutes using Amicon Ultra 15mL/100kD MWCO Centrifugal filters (EMD Millipore, Ref. No. UFC9100024). Total virus was then resuspended to a volume in pH 7.1 D10 media with polybrene so that each well would receive 500 μ L virus. To pH equilibrate media prior to infection, we combined virus with 600 μ L of pH 7.1 tissue culture media and incubated at 37°C and 5% CO₂ in a tissue culture incubator. We also aspirated all media from cells and added 1mL of the pH 7.1 tissue culture media. Both virus and cells were incubated separately for 60 minutes, after which 1mL of virus was added to the cells. Infection assays proceeded for ~12–14 hours.

For VSV-G expressing pseudovirus, we aimed for 100-fold coverage of barcoded variants and generally aimed to infect cells with ~12–15 million transducing units of virus. Due to this high number, we split VSV-G pseudovirus infections across two wells of 293T cells such that each well was infected with ~6 to 7.5 million transducing units. To ensure that infections occurred at an optimal pH for VSV-G fusion activity, we conducted these infections in tetracycline-negative standard D10 media (pH 7.4). VSV-G pseudovirus was also spin concentrated and resuspended in D10 media to a sufficient volume for each well of cells to receive 1 mL of virus. As for rabies G library pseudovirus, infections proceeded for 12–14 hours.

After the infection assays, cells were miniprep to collect non-integrated reverse-transcribed lentiviral genomes using a QIAprep Spin Miniprep Kit (Qiagen, Ref. No. 2710). To prepare cells for miniprepping, media was aspirated from infected cells, and cells were trypsinized (Fisher, Ref. No. MT25053C) and collected into microcentrifuge tubes. We washed cells with 1X PBS and miniprep the cell pellet. To increase DNA yield, we warmed miniprep columns with 35 μ L of elution buffer added to 55°C for 5 minutes prior to spin-down elution.

Deep mutational scanning for mutational effects on antibody escape

The experiments to measure the effects of G mutations on escape from antibody neutralization were performed largely as above with minor alterations. Firstly, to convert sequencing counts into absolute neutralization, we used a “neutralization standard” consisting of a barcoded pseudovirus expressing VSV-G as described previously.^{36,37} These pseudoviruses contain a defined set of barcodes and act as a non-neutralized standard for assays. This standard virus was added to the rabies G pseudovirus library at 1–2% of the rabies G pseudovirus library titer. As above, we used spin concentration to remove D10 media and to exchange with D10 media which has a pH of 7.1.

We performed infection assays at multiple concentrations of antibody to achieve levels of neutralization ranging from ~50% to >99% as determined by the sequencing and comparison to the neutralization standard as described below. We combined 500 μ L of resuspended virus with 600 μ L of media supplemented with antibody so that the combined 1.1mL mixture contained antibody at the appropriate concentrations. We allowed antibody binding and pH equilibration to occur for 1 hour at 37°C and 5% CO₂ in a tissue culture incubator as the cells in 1mL of pH 7.1 media. After this incubation period, virus was added to cells and infections

proceeded for 12–14 hours. Infected cells were washed and non-integrated lentiviral DNA was miniprep as above for cell entry experiments.

Illumina sequencing of pseudovirus barcodes

We used Illumina sequencing of the pseudovirus barcodes to quantify infection by each G variant in each condition. To prepare samples for sequencing, we employed a protocol involving two steps of PCR as described in Dadonaite et al.³⁶ We used the first round of PCR to amplify barcodes and to append Illumina Truseq Read 1 and Read 2 sequences. An Illumina Truseq Read 1 sequence was added via the forward primer annealing upstream of the barcode (Illumina_Rnd1_For), and the Read 2 sequence was added using the reverse primer annealing downstream of the barcode (Illumina_Rnd1_Rev). See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV-G_Sequencing-Primers.fasta for sequences of the Illumina primers.

For each PCR, we combined 25 or 28 μ L of miniprep non-integrated lentiviral DNA, 1.5 μ L of each primer (at 10 μ M stock), and 25 μ L of KOD Hot Start master mix. PCR cycling conditions were as follows:

95°C, 2 min
95°C, 20 sec
70°C, 1 sec
58°C, 10 s, cooling at 0.5°C/sec
70°C, 20 sec
Return to Step 2, 27 cycles
70°C, 1 min
4°C hold

We cleaned PCR products with 150 μ L of Ampure XP beads each (3:1 volume ratio of beads to PCR reaction) and eluted in 50 μ L of elution buffer. We quantified concentrations of the PCR product with a Qubit 4 Fluorometer (ThermoFisher, Ref. No. Q33238).

For the second step of the PCR, we appended the P5 Illumina adapter overhang using a forward primer that anneals to the Illumina Truseq Read 1 sequence (Illumina_Rnd2_For). The reverse primer annealed to the Truseq Read 2 sequence and contained the P7 Illumina adapter and i7 sample index (Illumina_Rnd2_Rev). Indices were from the NextFlex 8nt barcode set. See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV-G_Sequencing-Primers.fasta for sequences of the round 2 primers. Note that since indexes can vary, we have only provided a generic template where the index is denoted with 'n' characters. For each reaction, we combined 20 ng of round 1 PCR product, 2 μ L of each primer (at 10 μ M each) and added molecular biology grade water to 25 μ L total. To this, we added 25 μ L of KOD Hot Start Mix. We used PCR cycling conditions the same as above, except only 20 cycles total were performed.

We quantified round 2 PCR product concentration using a Qubit 4 Fluorometer (ThermoFisher, Ref. No. Q33238). We then pooled samples at masses (ng) proportional to the reads desired and ran the pool on a 1% agarose gel and extracted the band of the correct size (283bp). We ampure-cleaned the gel purified fragment and diluted the sequencing pool in elution buffer to a concentration of 4 nM. Prior to sequencing, we confirmed the final sample contained a monodisperse peak around the expected product size via TapeStation.

We aimed to have at least 20 million single-end Illumina reads per barcoded sample in a pooled submission. For VSV-G library pseudovirus, we aimed for over 50 million reads per sample to ensure adequate depth for calculating library composition.

Production of luciferase-encoding pseudovirus for validation assays

We generated pseudovirus expressing rabies G and encoding luciferase for validation assays of how specific G mutations affected cell entry or antibody neutralization. Briefly, we seeded 750,000 293T cells into each well of 6-well plates. For each well, we transfected 250 ng of G expression plasmid, 250 ng each of 26_HDM-Hgpm2 (Gagpol), 27_HDM-tat1b (Tat), and 28_pRC-CMV-Rev1b (Rev) expression plasmids, and 1 μ g of a plasmid containing a lentiviral genome encoding luciferase and driven by a chimeric CMV-LTR promoter. Virus was harvested and sterile filtered via syringe (Corning, Ref. No. 431220). We generated all viruses in standard D10 at a pH of 7.4.

To titrate these pseudoviruses, we seeded 15,000 293T cells per well in black-wall, clear-bottom, poly-L-Lysine coated 96-well plates 24 hours prior to infections. On the day of infection, media was removed and replaced with 50 μ L of fresh D10 at pH 7.1 supplemented with 4 μ g/mL polybrene. We generated serial dilution courses of virus in separate plates proceeding in 2- or 4- fold dilution steps. This allows us to sample RLU values at multiple volumes to gather technical replicates. Because the pH 7.1 D10 media requires pre-equilibration in tissue culture incubator settings, we pre-equilibrated virus and cells separately for 60 minutes. We then added 100 μ L to the cells for infections.

After 48 hours, we measured RLU using the Bright-Glo Luciferase Assay System (Promega, Ref. No. E2620). Briefly, 100 μ L of media was aspirated from wells, and 30 μ L of Bright-Glo was added for a 1:1 volume ratio in each well. Cells were lysed and luciferase activity was measured by plate-reader. To reduce background luminescence, we placed a black sticker surface to mask the clear bottoms of all wells. We normalized counts to volume to generate RLU/ μ L to calculate volume required for 250,000–500,000 RLU for neutralization curves. For each virus, we calculated an RLU/ μ L from technical replicates by taking the mean of all individual replicates.

Validation pseudovirus neutralization assays

For neutralization assays to validate the effects of specific G mutations, we seeded 15,000 293T cells per well in black-wall, clear-bottom, poly-L-Lysine coated 96-well plates 24 hours prior to infections. On the day of infection, media was removed and replaced with 50 μ L of fresh D10 at pH 7.1 supplemented with 4 μ g/mL polybrene. We generated an antibody dilution course plate using 5-fold dilutions steps down from a maximum concentration. Each plate had 2 column replicates for each antibody, as well as 2 column replicates for wells having no antibody to measure initial RLU per well. We also had a reserved column with no virus as an autofluorescence control.

We diluted the virus into D10 at pH 7.1 supplemented with 4 μ g/mL polybrene so that a volume of 60 μ L would yield approximately 250,000-500,000 RLU. To a new plate, we added 60 μ L to each well (except for those serving as a no-virus control). We combined 60 μ L from each well in the antibody plate and allowed binding and pH equilibration to proceed for 60 minutes. Following this, we took 100 μ L from each well and added them to the plate containing cells.

RLU for neutralization assays were measured as for luciferase titer assays. RLU values in each plate were mapped to a virus, antibody identity, and antibody concentration. We also took the median of luminescence values and subtracted each RLU measurement to correct for plate background. To calculate fraction infectivity, we calculated the median RLU of all wells containing no antibody and normalized all RLU measurements to this quantity. We fit neutralization curves to the fraction infectivities using the neutcurve package⁹⁰ (<https://jbloomlab.github.io/neutcurve/index.html>).

Pseudovirus validations of mutational effects on cell entry and antibody escape

To validate cell entry and antibody escape mutations in rabies G, we generated individually cloned luciferase viruses that express glycoprotein mutants. To do this, we first cloned point mutations in an expression plasmid encoding the Pasteur strain of rabies G (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/3968_HDM_Rabies-G_Pasteur_GenscriptOpt3.gb). Mutagenesis primers were generated using the NEBaseChanger tool. See (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PrimerSequences/RABV-G_Mutagenesis-Primers.fasta) for all of the mutagenesis primers. For each mutagenesis reaction, we combined 10ng of unmutated expression plasmid, 1.25 μ L each of corresponding forward and reverse primers (10 μ M stock), and 12.5 μ L of Q5 High-Fidelity 2X Master Mix (New England Biolabs, Ref. No. M0492S). We used the kit's standard PCR cycling conditions and suggested annealing temperatures. To digest unmutated template plasmid and ligate PCR products, the PCR products were treated with KLD Enzyme Mix (New England Biolabs, Ref. No. M0554S). We transformed a small volume of the PCR product into Stellar Chemically Competent cells (Takara, Ref. No. 636766). These expression plasmids were used as before to make luciferase-encoding pseudovirus (see “production of luciferase-encoding pseudovirus” section). See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/Additional_Data/PlasmidSequences/4954_HDM_Rabies-G_Pasteur_I338T.gb for an example mutant plasmid map.

We performed titrations using this virus as described above (see “Titration of virus via luciferase readout”). To compare with functional effects from the deep mutational scan, we normalized these titers to those of unmutated Pasteur rabies G pseudovirus (Figure 1D). For glycoprotein mutants with entry scores below -5 (H270P and N319A), we set cell entry to -5 as this was equivalent to the effect of stop codons. We performed replicates using independent stocks of miniprepmed expression plasmids. See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/non-pipeline_analyses/Additional_Notebooks/240810_LuciferaseValidations.ipynb for raw data and notebooks for analysis.

The validation assays for escape mutations from antibody 17C7 shown in Figure S3D were performed and analyzed as above in the “neutralization assays” section with minor changes. Firstly, we aimed for a maximum antibody concentration of 5 μ g/mL. Secondly, we adjusted the layout of the plates to accommodate the necessity of no antibody controls for 4 separate viruses per plate. See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/non-pipeline_analyses/Additional_Notebooks/17C7_EscapeValidations.ipynb for analysis.

Validation of deep mutational scanning predictions using natural rabies G strains

We identified rabies G protein sequences for two circulating strains A12_2718 (NCBI Accession No. KC792123.1) and NY-2011-12020 (NCBI Accession No. QJ74619.1) and codon sequence optimized using software from Twist or Genscript. We ordered these fragments from Twist and cloned them into the expression vector using HiFi Assembly. See https://github.com/dms-vep/RABV_Pasteur_G_DMS/tree/main/Additional_Data/PlasmidSequences for plasmid maps for the wild-type glycoproteins and for mutants generated by QuickChange (described below). The plasmid encoding the wild-type A12_2718 glycoprotein is 4956_HDM_Rabies-G_AGN94399-WT, and the plasmid encoding the wild-type NY-2011-12020 glycoprotein is 4959_HDM_Rabies-G_QJ74619_WT.

Strains were pseudotyped onto lentivirus and titered for luciferase activity as above with minor modifications. For producing pseudovirus expressing the NY-2011-12020 glycoprotein, we produced virus at a scale of ~60mL and concentrated via centrifugation into 1mL of virus to generate sufficient titers. We also performed plate-reader measurements in white-walled plates to reduce the amount of virus required.

To study the contribution of individual mutations in the A12_2718 background, we made mutations using the QuickChange protocol as described in the “pseudovirus validations of mutational effects on cell entry and antibody escape” section. Virus was rescued and concentrated for the T338I mutant glycoprotein. To ensure sufficient titers for neutralization assays, we also measured titers in white-wall plates. Neutralization assays for antibodies were performed as previously described in the “neutralization assays” section. With the exception of the experiment measuring neutralization of unmutated A12_2718 rabies G by 17C7 and RVC58

(shown in Figure 6C), we transferred the luciferase and cell mixture into white-wall plates for measurements for sufficient luciferase signal. Neutralization curves were subsequently processed using the neutcurve package as above (see “neutralization assays”).

Measuring cell surface expression levels of rabies G mutants

We measured cell surface expression levels of selected rabies G mutants (Figures S4B and S5E) by transfecting 293T cells with expression plasmids that encode rabies G mutants of interest. These plasmids were generated using the QuickChange protocol as described in “pseudovirus validations of mutational effects on cell entry and antibody escape.” We seeded 6-well plates with 750,000 293T cells each. The following day, we transfected each well with 500ng of a rabies G expression plasmid, supplemented with 1.5 μ g of transfection carrier DNA (Promega, Ref. No. E4881). These plasmids were combined with 200 μ L of DMEM and 3 μ L of BioT transfection reagent, and we let this mixture sit at room temperature for 15 minutes prior to adding to cells drop-wise. For our “No transfection” control, only cell culture media was present. Rabies G mutants were allowed to express for 48 hours in total. To prevent media pH drop that can lead to syncytia formation, we replaced D10 media in each well with fresh D10 media 24 hours after transfection.

After expression, media was aspirated by pipetting and cells were washed twice in 1X PBS solution while still bound to the plate. To preserve expressed rabies G proteins, cells were dissociated from plates using Cell Dissociation Buffer Enzyme-Free PBS-based (Gibco, Ref. No. 13151-014). Cells were then washed twice with 1X PBS supplemented with 3% Bovine Serum Albumin Fraction V (Gibco, Ref. No. 15260-037). We fixed cells in 1% paraformaldehyde solution (Biotium, Ref. No. 22023) and washed an additional two times with PBS and 3% BSA prior to staining with antibodies.

We labeled cells using pre-fusion specific rabies G antibodies as the primary antibodies, RVC20 (5.71mg/mL stock) and RVA122 (6.02mg/mL stock). Both antibodies were diluted 1:1000 each from their original stock in PBS with 3% BSA. Cells were incubated with primary antibody for 30 minutes, and then washed twice. We then incubated cells with a secondary antibody, Goat anti-Human IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647 (Invitrogen, Ref. No. A-21445) diluted 1:500 in PBS with 3% BSA, for 30 minutes. Cells were again washed twice. For flow cytometry, we used a Fortessa X50 instrument and detected staining using the APC filter. Flow cytometry data was analyzed using FlowJo and gated to eliminate debris and cell doublets. We plotted the histograms of labeling using the “modal” y-axis setting for Figures S4B and S5E.

Computational Methods

PacBio sequencing analysis

To analyze PacBio sequencing data to link the barcodes to the G mutant in the lentiviral backbones, we used the *alignparse* package (<https://github.com/jbloomlab/alignparse>).⁹¹ We subsequently filtered sequencing reads on various criteria. Firstly, we removed Circular Consensus Sequencing reads (CCSs) that contained an error rate higher than 10⁻⁴. This removed ~25% of reads for Libraries A and B. We also used the G or C nucleotide tags added during PCR to remove strand-exchange sequences, which removed ~1% of sequences per library (validating that the rate of strand exchange during the library-prep PCR was low). These CCSs were then aligned to a rabies G reference sequence, and we generated consensus sequences for each barcode requiring at least 3 CCSs and below 20% of minor sub or indel frequencies. We generated a barcode to variant look-up table with these sequences with an empirical accuracy of ~0.8 for both libraries. After this process, Library A contained 83,706 variants, and Library B contained 89,327 variants (Figure S1C).

See (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/results/variants/codon_variants.csv) for the final barcode to variant table. Jupyter notebooks are available with the code for analyzing PacBio CCSs (https://dms-vep.org/RABV_Pasteur_G_DMS/notebooks/analyze_pacbio_ccs.html), building consensus sequences (https://dms-vep.org/RABV_Pasteur_G_DMS/notebooks/build_pacbio_consensus.html) and building the final barcode-variant table (https://dms-vep.org/RABV_Pasteur_G_DMS/notebooks/build_codon_variants.html).

Illumina barcode sequencing analysis

We used the tool *illuminabarcodeparser* (https://jbloomlab.github.io/dms_variants/dms_variants.illuminabarcodeparser.html) to count barcodes mapping to each variant for each experimental conditions. We filtered for valid barcodes by examining nucleotide sequences downstream of a standard sequence in the amplicon with only 2 mismatches allowed. Barcodes were required to have a minimum quality score of 20. For each experiment, we required that barcodes met a minimum count threshold in “pre-selection” conditions, either VSV-G pseudovirus barcode counts in functional effect selections or no-antibody conditions in antibody escape assays. This step removes barcodes that are poorly represented in specific experiments. The specific thresholds used for each experiment is specified in the YAML configuration files on the GitHub repo (https://github.com/dms-vep/RABV_Pasteur_G_DMS).

Calculation of effects of mutations on cell entry

We first assigned a cell entry functional score to each barcoded variant.^{36,37} Briefly, for a barcode mutant m , we calculated functional scores using the formula $\log_2 [(n_{post}^m / n_{post}^{wt}) / (n_{pre}^m / n_{pre}^{wt})]$, where n_{post}^m is the barcode count for a mutant in the rabies G expressing Pseudovirus library, n_{pre}^m is the barcode count for a mutant in the VSV-G expressing library, n_{post}^{wt} and n_{pre}^{wt} are the corresponding counts for all unmutated (wildtype) G barcoded variants in the library. Therefore, G mutants that have wildtype-like cell entry will have a functional score of zero, while mutants with impaired G entry will have functional scores less than zero. Because some mutants in the library contain multiple amino acid mutations, we applied a global epistasis model⁴⁵ with a sigmoid global epistasis function as implemented in *multidms*⁹² to deconvolve the effects of individual amino acid mutations. In this paper and in the interactive displays at https://dms-vep.org/RABV_Pasteur_G_DMS/cell_entry.html we report the average cell entry effects for mutations across libraries

and experiments. Basic filtering for high-confidence measurements was applied by only reporting values for mutations seen in an average of at least two barcoded variants per library and with measurements in the majority of experimental replicates.

Calculation of effects of mutations on antibody escape

To calculate the effects of mutations on antibody escape, for each barcoded variant, we first determined the non-neutralized fraction at each antibody concentration by comparing the barcode counts for the variant to those for the non-neutralized VSV-G standard barcodes as previously described in Dadonaite et al.³⁶ We then used the *polyclonal*⁵¹ software package (<https://jbloomlab.github.io/polyclonal/>) to estimate the effects of each mutation on escape. The mutation effects on escape are linearly correlated with the change in log IC50 caused by each mutation, with positive values indicating escape and negative values indicating increased neutralization. See the “antibody/serum escape” section of https://dms-vep.org/RABV_Pasteur_G_DMS/appendix.html for notebooks with details about the fitting and that show the fraction of the library that is non-neutralized at each antibody concentration.

For all figures and interactive visualizations (https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html), we applied filters to remove low-confidence measurements. Specifically, we only showed mutations observed in an average of at least two-barcoded variants across libraries, and by default we hide the escape effects for mutations that are highly deleterious for cell entry (as it is difficult to measure the neutralization of G variants that are largely non-functional). The default visualizations also floor escape values at zero (so only show positive escape). All of these options can be adjusted using the interactive visualizations at https://dms-vep.org/RABV_Pasteur_G_DMS/escape.html.

Nextstrain phylogenetic trees of publicly available rabies G sequences

We gathered publicly available rabies G sequences from NCBI Virus using the query term “Lyssavirus rabies, taxid:11292” in the “Search by virus name or taxonomy box.” Sequences were downloaded as of October 18, 2024. Only G sequences with at least 1500 nucleotides and no ambiguous nucleotides (“N”) were included. The G sequences were processed and aligned using Augur,¹⁰¹ and a phylogenetic tree was inferred from the resulting alignment. The phylogenetic tree was rooted using the non-rabies lyssavirus (also in phylogroup I) Gannoruwa bat lyssavirus (NCBI accession: NC_031988) as an outgroup.

Antibody escape scores were mapped onto the tree by summing the escape measurements of G sequence substitutions relative to the Pasteur G sequence, and the tree was visualized using Nextstrain.⁷⁸ See <https://nextstrain.org/groups/jbloomlab/dms/rabies-G> for the interactive Nextstrain trees, and https://github.com/dms-vep/RABV_Pasteur_G_DMS/tree/main/non-pipeline_analyses/RABV_nextstrain for the computer code used to generate the trees. Note that the trees can be colored by G amino-acid identity using either the G ectodomain numbering scheme used in this paper, or sequential numbering of the entire G protein.

Note that the Nextstrain phylogenetic trees use all 7,122 publicly available sequences, but the plot in Figure 6A uses only the 6,581 sequences corresponding to natural sequences (excluding lab-passaged ones and those originating from an unknown host).

Structural analysis

We examined the effects of mutations on several previously determined structures of rabies G. These structures include pre-fusion conformation (PDB: 7U9G), extended intermediate conformation (PDB: 6LGW) and rabies G in complex with antibodies (PDB 8R40 for CR57, PDB 8A1E for 17C7, PDB 7U9G for RVA122, and PDB 6TOU for RVC20). To project the mutation effects on structures, we made a notebook (https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/non-pipeline_analyses/Additional_Notebooks/color_pdb_functionaleffects.ipynb) that takes a PDB file, relevant chain IDs, and a CSV of deep mutational scan measurements and re-assigns the b-factors in the PDB file to the desired measurements. For cell-entry we used a site statistic of the mean effect of mutations at each site; for antibody escape we used the summed positive escape values across all mutations at that site.

See https://github.com/dms-vep/RABV_Pasteur_G_DMS/blob/main/non-pipeline_analyses/Additional_Notebooks/241126_GetContacts.ipynb for jupyter notebooks used to determine all residues within 4Å of an antibody in complex structures and producing heatmaps.

QUANTIFICATION AND STATISTICAL ANALYSIS

All code and quantitative analysis is publicly available on GitHub (https://github.com/dms-vep/RABV_Pasteur_G_DMS). Interactive plots which show metrics for each measurement are publicly available an easy-to-view HTML summary (https://dms-vep.org/RABV_Pasteur_G_DMS/). Functional effects on cell entry were calculated as defined in “[calculation of effects of mutations on cell entry](#)” of Methods using the *multidms* software package⁹² (<https://matsengrp.github.io/multidms/>) to implement a global epistasis model with a sigmoid function.⁴⁵ Antibody escape was calculated using the *polyclonal* software package⁵¹ (<https://jbloomlab.github.io/polyclonal/>) as described in “[calculation of effects of mutations on antibody escape](#)” of Methods. Neutralization curves were generated using *neutcurve*⁹⁰ (<https://jbloomlab.github.io/neutcurve/index.html>). IC50 values were calculated using the software’s standard protocol of averaging technical duplicates of each measurement in a neutralization curve.

Summary statistics for measurements are reported directly in figure legends. For heatmaps, we report the mean of all replicates; mutants reported were also required to have a *times_seen* value of at least 2 in each replicate library. We projected the mutational effects on cell entry and antibody escape on structures as defined in the “[structural analysis](#)” section of Methods and as listed in the figure legends.